arXiv:1502.02423v3 [math.QA] 15 Mar 2015

ON THE MINIMAL AFFINIZATIONS OF TYPE F_4

BING DUAN, JIAN-RONG LI, YAN-FENG LUO

Abstract. In this paper, we apply the theory of cluster algebras to study minimal affinizations over the quantum affine algebra of type F_4. We show that the q-characters of a large family of minimal affinizations of type F_4 satisfy some equations. Moreover, every minimal affinization these equations corresponds to some cluster variable in some cluster algebra $\mathscr{A}$. For the other minimal affinizations of type F_4 which are not these equations, we give some conjectural equations which contains these minimal affinizations. Furthermore, we introduce the concept of dominant monomial graphs to study the equations satisfied by q-characters of modules of quantum affine algebras.

Key words: quantum affine algebra of type F_4; minimal affinizations; cluster algebras; q-characters; Frenkel-Mukhin algorithm

2010 Mathematics Subject Classification: 17B37; 13B60

1. Introduction

The theory of cluster algebras are introduced by Fomin and Zelevinsky in [FZ02]. It has many applications to mathematics and physics including quiver representations, Teichmüller theory, tropical geometry, integrable systems, and Poisson geometry.

Let $\mathfrak{g}$ be a simple Lie algebra and $U_q\widehat{\mathfrak{g}}$ the corresponding quantum affine algebra. In [C95], V. Chari and A. Pressley introduced minimal affinizations of representations of quantum groups. The family of minimal affinizations is an important family of simple modules which contains the Kirillov-Reshetikhin modules. Minimal affinizations are studied intensively in recent years, see for examples, [CG11], [CMY13], [Her07], [Li15], [LM13], [M10], [MP07], [MP11], [MY12a], [MY12b], [MY14], [Nao13], [Nao14], [QL14], [SS14], [ZDLL15]. However, there is not much work of the minimal affinizations over the quantum affine algebra of type F_4 in the literature. The aim of this paper is to apply the theory of cluster algebras to study minimal affinizations over the quantum affine algebra of type F_4.

M-systems and dual M-systems of types A_n, B_n, G_2 are introduced in [ZDLL15], [QL14] to study the minimal affinizations of types A_n, B_n, G_2. The equations in these systems are satisfied by the q-characters of minimal affinizations of types A_n, B_n, G_2. It is shown that every equation in these systems corresponds to a mutation equation in some cluster algebra.

The minimal affinizations of type F_4 are much more complicated than the minimal affinizations of types A_n, B_n, G_2. In types A_n, B_n, G_2, all minimal affinizations are special or anti-special, [Her07], [LM13]. A $U_q\widehat{\mathfrak{g}}$-module V is called special (resp. anti-special) if there is only one dominant (resp. anti-dominant) monomial in the q-character of V. In the case of type F_4, there are minimal affinizations which are neither special nor anti-special, [Her07].

In [ZDLL15], [QL14], the M-systems and dual M-systems of types A_n, B_n, G_2 contain all minimal affinizations and only contain minimal affinizations. The situation is different in the case of type F_4. It is quite possible that a closed system which contains all minimal affinizations and only contains minimal affinizations of type F_4 does not exist. However, we are able to find two closed systems which contain a large family of minimal affinizations of type F_4, Theorem 3.4, Theorem 5.4. We show that the equations these systems are satisfied by the q-characters of the minimal affinizations in the systems. We prove that the modules in one system are special, Theorem 3.3, and the modules in the other system are anti-special, Theorem 5.2. Moreover, we show that every equation in Theorem 3.4 (resp. Theorem 5.4) corresponds to a mutation equation in some cluster algebra $\mathscr{A}$ (resp. $\widetilde{\mathscr{A}}$). The cluster algebra $\mathscr{A}$ is the same as the cluster algebra for the quantum affine algebra of type F_4 introduced in [HL13]. Moreover, every minimal affinization in Theorem 3.4 (resp. Theorem 5.4) corresponds to a cluster variable in the cluster algebra $\mathscr{A}$ (resp. $\widetilde{\mathscr{A}}$), Theorem 4.1 (resp. Theorem 5.6).

The system of equations in Theorem 5.4 is dual to the system in Theorem 3.4. This system contains the modules which are dual to the modules in the system in Theorem 3.4.

For the minimal affinizations which are not in Theorem 3.4 and Theorem 5.4, we give some conjectural equations which contain these modules, Conjecture 9.1.

We introduce the concept of dominant monomial graphs to study the equations satisfied by q-characters of modules of quantum affine algebras. We draw dominant monomial graphs for the modules in the equivalence classes of the left hand side of some equations in Conjecture 9.1. In these graphs, we find that every graph can be divided into two parts. The vertices in the first (resp. second) part of the graph are dominant monomials in the first (resp. second) summand of the right hand side of the correpsonding equation.

The paper is organized as follows. In Section 2, we give some background information about cluster algebras and representation theory of quantum affine algebras. In Section 3, we describe a closed system containing a large family of minimal affinizations of type F_4. In Section 4, we study relations between the system in Theorem 3.4 and cluster algebras. In Section 5, we study the dual system of Theorem 3.4. In Section 6, Section 7, and Section 8, we prove Theorem 3.3, Theorem 3.4, and Theorem 3.6 given in Section 3, respectively. In Section 9, we give a conjecture about the equations satisfied by the q-characters of the other minimal affinizations of type F_4 and introduce the concept of dominant monomial graphs to study the conjecture.

2. BACKGROUND

2.1. **Cluster algebras.** We first recall the definition of cluster algebras introduced by Fomin and Zelevinsky in [FZ02]. Let $\mathbb{Q}$ be the rational field and $\mathcal{F} = \mathbb{Q}(x_1, x_2, \cdots, x_n)$ the field of rational functions in n indeterminates over $\mathbb{Q}$. A seed in $\mathcal{F}$ is a pair $\Sigma = (\mathbf{y}, Q)$, where $\mathbf{y} = (y_1, y_2, \cdots, y_n)$ is a free generating set of $\mathcal{F}$, and Q is a quiver with vertices labeled by $\{1, 2, \cdots, n\}$. Assume that Q has neither loops nor 2-cycles. For $k \in \{1, 2, \cdots, n\}$, one defines a new seed $\mu_k(\mathbf{y}, Q) = (\mathbf{y}', Q')$ by the mutation of $(\mathbf{y}, Q)$ at k. Here $\mathbf{y}' = (y_1', \ldots, y_n')$, $y_i' = y_i$, for $i \neq k$, and

$$y_k' = \frac{\prod_{i \to k} y_i + \prod_{k \to j} y_j}{y_k}, \tag{2.1}$$

where the first (resp. second) product in the right hand side is over all arrows of Q with target (resp. source) k, and Q' is obtained from Q by the follow rule:

(i) Reverse the orientations of all arrow incident with k;
(ii) Add a new arrow $i \to j$ for every existing pair of arrow $i \to k$ and $k \to j$;

(iii) Erasing every pair of opposite arrows possible created by (ii).

The mutation class $\mathcal{C}(\Sigma)$ is the set of all seeds obtained from Σ by a finite sequence of mutation μ_k. If $\Sigma' = ((y_1', y_2', \cdots, y_n'), Q')$ is a seed in $\mathcal{C}(\Sigma)$, then the subset $\{y_1', y_2', \cdots, y_n'\}$ is called a *cluster*, and its elements are called *cluster variables.* The *cluster algebra* $\mathscr{A}_\Sigma$ as the subring of $\mathcal{F}$ generated by all cluster variables. *Cluster monomials* are monomials in the cluster variables supported on a single cluster.

In this paper, the initial seed in the cluster algebra we use is of the form $\Sigma = (\mathbf{y}, Q)$, where $\mathbf{y}$ is an infinite set and Q is an infinite quiver.

Definition 2.1 (Definition 3.1, [GG14])**.** *Let Q be a quiver without loops or 2-cycles and with a countably infinite number of vertices labelled by all integers $i \in \mathbb{Z}$. Furthermore, for each vertex i of Q let the number of arrows incident with i be finite. Let $\mathbf{y} = \{y_i \mid i \in \mathbb{Z}\}$. An infinite initial seed is the pair $(\mathbf{y}, Q)$. By finite sequences of mutation at vertices of Q and simultaneous mutation of the set $\mathbf{y}$ using the exchange relation (2.1), one obtains a family of infinite seeds. The sets of variables in these seeds are called the infinite clusters and their elements are called the cluster variables. The cluster algebra of infinite rank of type Q is the subalgebra of $\mathbb{Q}(\mathbf{y})$ generated by the cluster variables.*

2.2. **The quantum affine algebra of type F_4.** In this paper, we take $\mathfrak{g}$ to be the complex simple Lie algebra of type F_4 and $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$. Let $I = \{1, 2, 3, 4\}$. We choose simple roots $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ and scalar product $(\cdot, \cdot)$ such that

$$\begin{aligned}&(\alpha_1, \alpha_1) = 2,\ (\alpha_1, \alpha_2) = -1,\ (\alpha_2, \alpha_2) = 2,\ (\alpha_2, \alpha_3) = -2,\\ &(\alpha_3, \alpha_3) = 4,\ (\alpha_3, \alpha_4) = -2,\ (\alpha_4, \alpha_4) = 4.\end{aligned}$$

Therefore α_1, α_2 are the short simple roots and α_3, α_4 are the long simple roots.

Let $\{\alpha_1^\vee, \alpha_2^\vee, \alpha_3^\vee, \alpha_4^\vee\}$ and $\{\omega_1, \omega_2, \omega_3, \omega_4\}$ be the sets of simple coroots and fundamental weights respectively. Let $C = (C_{ij})_{i,j \in I}$ denote the Cartan matrix, where $C_{ij} = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$. Let $d_1 = 1, d_2 = 1, d_3 = 2, d_4 = 2$, $D = \mathrm{diag}(d_1, d_2, d_3, d_4)$ and $B = DC = (b_{ij})_{i,j \in I}$. Then

$$C = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -2 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -2 & 0 \\ 0 & -2 & 4 & -2 \\ 0 & 0 & -2 & 4 \end{pmatrix}.$$

Let $q_i = q^{d_i}$, $i \in I$. Let Q (resp. Q^+) and P (resp. P^+) denote the $\mathbb{Z}$-span (resp. $\mathbb{Z}_{\geq 0}$-span) of the simple roots and fundamental weights respectively. Let $\leq$ be the partial order on P in which $\lambda \leq \lambda'$ if and only if $\lambda' - \lambda \in Q^+$.

Quantum groups are introduced independently by Jimbo [Jim85] and Drinfeld [Dri87]. Quantum affine algebras form a family of infinite-dimensional quantum groups. Let $\widehat{\mathfrak{g}}$ denote the untwisted affine algebra corresponding to $\mathfrak{g}$. In this paper, we fix a $q \in \mathbb{C}^\times$, not a root of unity. The quantum affine algebra $U_q\widehat{\mathfrak{g}}$ in Drinfeld's new realization, see [Dri88], is generated by $x_{i,n}^{\pm}$ $(i \in I, n \in \mathbb{Z})$, $k_i^{\pm 1}$ $(i \in I)$, $h_{i,n}$ $(i \in I, n \in \mathbb{Z}\backslash\{0\})$ and central elements $c^{\pm 1/2}$, subject to certain relations.

The algebra $U_q\mathfrak{g}$ is isomorphic to a subalgebra of $U_q\widehat{\mathfrak{g}}$. Therefore $U_q\widehat{\mathfrak{g}}$-modules restrict to $U_q\mathfrak{g}$-modules.

2.3. **Finite-dimensional $U_q\widehat{\mathfrak{g}}$-modules and their q-characters.** We recall some known results on finite-dimensional $U_q\widehat{\mathfrak{g}}$-modules and their q-characters, [CP94], [CP95a], [FR98], [MY12a].

Let $\mathcal{P}$ be the free abelian multiplicative group of monomials in infinitely many formal variables $(Y_{i,a})_{i\in I, a\in\mathbb{C}^\times}$. Then $\mathbb{Z}\mathcal{P} = \mathbb{Z}[Y_{i,a}^{\pm 1}]_{i\in I, a\in\mathbb{C}^\times}$. For each $j \in I$, a monomial $m = \prod_{i\in I, a\in\mathbb{C}^\times} Y_{i,a}^{u_{i,a}}$, where $u_{i,a}$ are some integers, is said to be *j-dominant* (resp. *j-anti-dominant*) if and only if $u_{j,a} \geq 0$ (resp. $u_{j,a} \leq 0$) for all $a \in \mathbb{C}^\times$. A monomial is called *dominant* (resp. *anti-dominant*) if and only if it is j-dominant (resp. j-anti-dominant) for all $j \in I$.

Every finite-dimensional simple $U_q\widehat{\mathfrak{g}}$-module is parametrized by a dominant monomial in $\mathcal{P}^+$, [CP94], [CP95a]. That is, for a dominant monomial $m = \prod_{i\in I, a\in\mathbb{C}^\times} Y_{i,a}^{u_{i,a}}$, there is a corresponding simple $U_q\widehat{\mathfrak{g}}$-module $L(m)$. Let $\mathrm{Rep}(U_q\widehat{\mathfrak{g}})$ be the Grothendieck ring of finite-dimensional $U_q\widehat{\mathfrak{g}}$-modules and $[V] \in \mathrm{Rep}(U_q\widehat{\mathfrak{g}})$ the class of a finite-dimensional $U_q\widehat{\mathfrak{g}}$-module V.

The q-character of a $U_q\widehat{\mathfrak{g}}$-module V is given by

$$\chi_q(V) = \sum_{m\in\mathcal{P}} \dim(V_m) m \in \mathbb{Z}\mathcal{P},$$

where V_m is the l-weight space with l-weight m, see [FR98]. We use $\mathscr{M}(V)$ to denote the set of all monomials in $\chi_q(V)$ for a finite-dimensional $U_q\widehat{\mathfrak{g}}$-module V. Let $\mathcal{P}^+ \subset \mathcal{P}$ denote the set of all dominant monomials. For $m_+ \in \mathcal{P}^+$, we use $\chi_q(m_+)$ to denote $\chi_q(L(m_+))$. We also write $m \in \chi_q(m_+)$ if $m \in \mathscr{M}(\chi_q(m_+))$.

The following lemma is well-known.

Lemma 2.2. *Let m_1, m_2 be two monomials. Then $L(m_1m_2)$ is a sub-quotient of $L(m_1)\otimes L(m_2)$. In particular, $\mathscr{M}(L(m_1m_2)) \subseteq \mathscr{M}(L(m_1))\mathscr{M}(L(m_2))$.* □

A finite-dimensional $U_q\widehat{\mathfrak{g}}$-module V is said to be *special* if and only if $\mathscr{M}(V)$ contains exactly one dominant monomial. It is called *anti-special* if and only if $\mathscr{M}(V)$ contains exactly one anti-dominant monomial. Clearly, if a module is special or anti-special, then it is simple.

Let $a \in \mathbb{C}^\times$ and

$$\begin{aligned} A_{1,a} &= Y_{1,aq}Y_{1,aq^{-1}}Y_{2,a}^{-1}, \\ A_{2,a} &= Y_{2,aq^1}Y_{2,aq^{-1}}Y_{1,a}^{-1}Y_{3,a}^{-1}, \\ A_{3,a} &= Y_{3,aq^2}Y_{3,aq^{-2}}Y_{4,a}^{-1}Y_{2,aq^1}^{-1}Y_{2,aq^{-1}}^{-1}, \\ A_{4,a} &= Y_{4,aq^2}Y_{4,aq^{-2}}Y_{3,a}^{-1}. \end{aligned} \tag{2.2}$$

Let $\mathcal{Q}$ be the subgroup of $\mathcal{P}$ generated by $A_{i,a}, i \in I, a \in \mathbb{C}^\times$. Let $\mathcal{Q}^\pm$ be the monoids generated by $A_{i,a}^{\pm 1}, i \in I, a \in \mathbb{C}^\times$. There is a partial order $\leq$ on $\mathcal{P}$ in which

$$m \leq m' \text{ if and only if } m'm^{-1} \in \mathcal{Q}^+. \tag{2.3}$$

For all $m_+ \in \mathcal{P}^+$, $\mathscr{M}(L(m_+)) \subset m_+\mathcal{Q}^-$, see [FM01].

The concept of *right negative* is introduced in Section 6 of [FM01].

Definition 2.3. *A monomial m is called right negative if for all $a \in \mathbb{C}^\times$, for $L = \max\{l \in \mathbb{Z} \mid u_{i,aq^l}(m) \neq 0$ for some $i \in I\}$ we have $u_{j,aq^L}(m) \leq 0$ for $j \in I$.*

For $i \in I, a \in \mathbb{C}^\times$, $A_{i,a}^{-1}$ is right-negative. A product of right-negative monomials is right-negative. If m is right-negative and $m' \leq m$, then m' is right-negative, see [FM01], [Her06].

2.4. **q-characters of $U_q\widehat{\mathfrak{sl}}_2$-modules and the Frenkel-Mukhin algorithm.** We recall the results of the q-characters of $U_q\widehat{\mathfrak{sl}}_2$-modules which are well-understood, see [CP91], [FR98].

Let $W_k^{(a)}$ be the irreducible representation $U_q\widehat{\mathfrak{sl}}_2$ with highest weight monomial

$$X_k^{(a)} = \prod_{i=0}^{k-1} Y_{aq^{k-2i-1}},$$

where $Y_a = Y_{1,a}$. Then the q-character of $W_k^{(a)}$ is given by

$$\chi_q(W_k^{(a)}) = X_k^{(a)} \sum_{i=0}^{k} \prod_{j=0}^{i-1} A^{-1}_{aq^{k-2j}},$$

where $A_a = Y_{aq^{-1}}Y_{aq}$.

For $a \in \mathbb{C}^\times, k \in \mathbb{Z}_{\geq 1}$, the set $\Sigma_k^{(a)} = \{aq^{k-2i-1}\}_{i=0,\dots,k-1}$ is called a *string*. Two strings $\Sigma_k^{(a)}$ and $\Sigma_{k'}^{(a')}$ are said to be in *general position* if the union $\Sigma_k^{(a)} \cup \Sigma_{k'}^{(a')}$ is not a string or $\Sigma_k^{(a)} \subset \Sigma_{k'}^{(a')}$ or $\Sigma_{k'}^{(a')} \subset \Sigma_k^{(a)}$.

Denote by $L(m_+)$ the irreducible $U_q\widehat{\mathfrak{sl}}_2$-module with highest weight monomial m_+. Let $m_+ \neq 1$ and $\in \mathbb{Z}[Y_a]_{a\in\mathbb{C}^\times}$ be a dominant monomial. Then m_+ can be uniquely (up to permutation) written in the form

$$m_+ = \prod_{i=1}^{s} \left(\prod_{b\in\Sigma_{k_i}^{(a_i)}} Y_b \right),$$

where s is an integer, $\Sigma_{k_i}^{(a_i)}, i = 1, \dots, s$, are strings which are pairwise in general position and

$$L(m_+) = \bigotimes_{i=1}^{s} W_{k_i}^{(a_i)}, \qquad \chi_q(L(m_+)) = \prod_{i=1}^{s} \chi_q(W_{k_i}^{(a_i)}).$$

For $j \in I$, let

$$\beta_j : \mathbb{Z}[Y_{i,a}^{\pm 1}]_{i\in I; a\in\mathbb{C}^\times} \to \mathbb{Z}[Y_a^{\pm 1}]_{a\in\mathbb{C}^\times}$$

be the ring homomorphism such that for all $a \in \mathbb{C}^\times$, $Y_{k,a} \mapsto 1$ for $k \neq j$ and $Y_{j,a} \mapsto Y_a$.

Let V be a $U_q\widehat{\mathfrak{g}}$-module. Then $\beta_i(\chi_q(V))$, $i \in I$, is the q-character of V considered as a $U_{q_i}\widehat{\mathfrak{sl}_2}$-module.

The Frenkel-Mukhin algorithm is introduced in Section 5 in [FM01] to compute the q-characters of $U_q\widehat{\mathfrak{g}}$-modules. In Theorem 5.9 of [FM01], it is shown that the Frenkel-Mukhin algorithm works for modules which are special.

2.5. **Truncated q-characters.** In this paper, we need to use the concept truncated q-characters, see [HL10], [MY12a]. Given a set of monomials $\mathcal{R} \subset \mathcal{P}$, let $\mathbb{Z}\mathcal{R} \subset \mathbb{Z}\mathcal{P}$ denote the $\mathbb{Z}$-module of formal linear combinations of elements of $\mathcal{R}$ with integer coefficients. Define

$$\mathrm{trunc}_{\mathcal{R}} : \mathcal{P} \to \mathcal{R}; \quad m \mapsto \begin{cases} m & \text{if } m \in \mathcal{R}, \\ 0 & \text{if } m \notin \mathcal{R}, \end{cases}$$

and extend $\mathrm{trunc}_{\mathcal{R}}$ as a $\mathbb{Z}$-module map $\mathbb{Z}\mathcal{P} \to \mathbb{Z}\mathcal{R}$.

Given a subset $U \subset I \times \mathbb{C}^\times$, let $\mathcal{Q}_U$ be the subgroups of $\mathcal{Q}$ generated by $A_{i,a}$ with $(i,a) \in U$. Let $\mathcal{Q}_U^{\pm}$ be the monoid generated by $A_{i,a}^{\pm 1}$ with $(i,a) \in U$. The polynomial $\mathrm{trunc}_{m_+\mathcal{Q}_U^-}\, \chi_q(m_+)$ is called *the q-character of $L(m_+)$ truncated to U.*

The following theorem can be used to compute some truncated q-characters.

Theorem 2.4 ([MY12a] , Theorem 2.1)**.** *Let $U \subset I \times \mathbb{C}^\times$ and $m_+ \in \mathcal{P}^+$. Suppose that $\mathcal{M} \subset \mathcal{P}$ is a finite set of distinct monomials such that*

(i) $\mathscr{M} \subset m_+\mathcal{Q}_U^-$,
(ii) $\mathcal{P}^+ \cap \mathscr{M} = \{m_+\}$,
(iii) for all $m \in \mathscr{M}$ and all $(i,a) \in U$, if $mA_{i,a}^{-1} \notin \mathscr{M}$, then $mA_{i,a}^{-1}A_{j,b} \notin \mathscr{M}$ unless $(j,b) = (i,a)$,
(iv) for all $m \in \mathscr{M}$ and all $i \in I$, there exists a unique i-dominant monomial $M \in \mathscr{M}$ such that

$$\mathit{trunc}_{\beta_i(M\mathcal{Q}_U^-)}\, \chi_q(\beta_i(M)) = \sum_{m' \in m\mathcal{Q}_{\{i\}\times\mathbb{C}^\times} \cap \mathscr{M}} \beta_i(m').$$

Then

$$\mathit{trunc}_{m_+\mathcal{Q}_U^-}\, \chi_q(m_+) = \sum_{m \in \mathscr{M}} m.$$

Here $\chi_q(\beta_i(M))$ is the q-character of the irreducible $U_{q_i}(\hat{\mathfrak{sl}}_2)$-module with highest weight monomial $\beta_i(M)$ and $\mathrm{trunc}_{\beta_i(M\mathcal{Q}_U^-)}$ is the polynomial obtained from $\chi_q(\beta_i(M))$ by keeping only the monomials of $\chi_q(\beta_i(M))$ in the set $\beta_i(M\mathcal{Q}_U^-)$.

2.6. **Minimal affinizations of $U_q\mathfrak{g}$-modules.** In what follows, we fix an $a \in \mathbb{C}^\times$ and denote $i_s = Y_{i,aq^s}$, $i \in I$, $s \in \mathbb{Z}$. Let $\lambda = k\omega_1 + l\omega_2 + m\omega_3 + n\omega_4$, $k,l,m,n \in \mathbb{Z}_{\geq 0}$ and $V(\lambda)$ the simple $U_q\mathfrak{g}$-module with highest weight λ. Without loss of generality, we may assume that a simple $U_q\widehat{\mathfrak{g}}$-module $L(m_+)$ is a *minimal affinization* of $V(\lambda)$ if and only if m_+ is one of the following monomials:

$$\widetilde{T}^{(s)}_{n,m,l,k} = \left(\prod_{i=0}^{n-1} Y_{4,aq^{s+4i}}\right)\left(\prod_{i=0}^{m-1} Y_{3,aq^{s+4n+4i+2}}\right)\left(\prod_{i=0}^{l-1} Y_{2,aq^{s+4n+4m+2i+3}}\right)\left(\prod_{i=0}^{k-1} Y_{1,aq^{s+4n+4m+2l+2i+4}}\right),$$

$$T^{(s)}_{n,m,l,k} = \left(\prod_{i=0}^{n-1} Y_{4,aq^{-s-4i}}\right)\left(\prod_{i=0}^{m-1} Y_{3,aq^{-s-4n-4i-2}}\right)\left(\prod_{i=0}^{l-1} Y_{2,aq^{-s-4n-4m-2i-3}}\right)\left(\prod_{i=0}^{k-1} Y_{1,aq^{-s-4n-4m-2l-2i-4}}\right),$$

where $\prod_{0\leq j\leq -1} = 1$, $s \in \mathbb{Z}$, see [CP96a]. We denote A^{-1}_{i,aq^s} by $A^{-1}_{i,s}$. We use $\mathcal{T}^{(s)}_{k,l,m,n}$ (resp. $\widetilde{\mathcal{T}}^{(s)}_{k,l,m,n}$) to denote the irreducible finite-dimensional $U_q\widehat{\mathfrak{g}}$-module with highest l-weight $T^{(s)}_{k,l,m,n}$ (resp. $\widetilde{T}^{(s)}_{k,l,m,n}$).

3. A CLOSED SYSTEM WHICH CONTAINS A LARGE FAMILY OF MINIMAL AFFINIZATIONS OF TYPE F_4

In this section, we introduce a closed system of type F_4 that contains a large family of minimal affinizations:

$$\begin{gathered}\mathcal{T}^{(-2)}_{0,0,l,k},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{0,m,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,0,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,0,k},\ \widetilde{\mathcal{T}}^{(s)}_{0,m,0,k}(k \leq 2),\\ \widetilde{\mathcal{T}}^{(s)}_{n,m,0,k}(k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}(k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{n,0,l,k}(k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{0,m,l,k}(k \leq 2).\end{gathered}$$

Here $k, l, m, n \in \mathbb{Z}_{\geq 1}$, $s \in \mathbb{Z}$.

3.1. **Special modules.**

Let $k, l, m, n \in \mathbb{Z}_{\geq 1}$, $s \in \mathbb{Z}$. For $k \leq 0$, let $T^{(s)}_{0,0,0,k} = 1$. We define

$$\begin{aligned}\widetilde{P}^{(s)}_{0,0,l,k} &= \widetilde{T}^{(s)}_{0,0,l,k}, \quad k \in \mathbb{Z}_{\leq 2},\\ \widetilde{P}^{(s)}_{0,m,0,k} &= \widetilde{T}^{(s)}_{0,m,0,k}, \quad k \in \mathbb{Z}_{\leq 2},\\ \widetilde{P}^{(s)}_{0,0,0,k} &= \widetilde{T}^{(s)}_{0,0,0,k}, \quad k \in \mathbb{Z}_{\geq 3},\\ \widetilde{P}^{(s)}_{0,0,l,k} &= \widetilde{T}^{(s+2l+4)}_{0,0,0,k-2}\widetilde{T}^{(s+2l+8)}_{0,0,0,k-4}\widetilde{T}^{(s)}_{0,0,l,k}, \quad k \in \mathbb{Z}_{\geq 3},\\ \widetilde{P}^{(s)}_{0,m,0,k} &= \widetilde{T}^{(s)}_{0,m,0,k}\widetilde{T}^{(s+4m+4)}_{0,0,0,k-2}, \quad k \in \mathbb{Z}_{\geq 3}.\end{aligned}$$

We use $\widetilde{\mathcal{P}}^{(s)}_{n,m,l,k}$ to denote the simple $U_q\widehat{\mathfrak{g}}$-module with the highest weight monomial $\widetilde{P}^{(s)}_{n,m,l,k}$.

Theorem 3.1 (Theorem 3.9, [Her07]). *For $l, m, n \in \mathbb{Z}_{\geq 1}$, $s \in \mathbb{Z}$, the modules $\widetilde{\mathcal{T}}^{(s)}_{n,m,0,0}$, $\widetilde{\mathcal{T}}^{(s)}_{n,0,l,0}$, $\widetilde{\mathcal{T}}^{(s)}_{0,m,l,0}$, $\widetilde{\mathcal{T}}^{(s)}_{n,m,l,0}$ are special.*

Remark 3.2. *In the paper* [Her07], *α_1, α_2 are simple long roots and α_3, α_4 are simple short roots. In this paper, α_1, α_2 are simple short roots and α_3, α_4 are simple long roots.*

Theorem 3.3. *The modules*

$$\begin{gathered}\mathcal{T}^{(-2)}_{0,0,l,k},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{0,m,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,0,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,0,k}, \widetilde{\mathcal{T}}^{(s)}_{n,m,0,k}\ (k \leq 2),\\ \widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}\ (k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{n,0,l,k}\ (k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{0,m,l,k}\ (k \leq 2),\ \widetilde{P}^{(s)}_{0,0,l,k},\ \widetilde{P}^{(s)}_{0,m,0,k},\end{gathered}$$

where $k, l, m, n \in \mathbb{Z}_{\geq 1}$, $s \in \mathbb{Z}$, are special.

We will prove Theorem 3.3 in Section 6.

Since the modules are special in Theorem 3.1 and Theorem 3.3, we can use the Frenkel-Mukhin algorithm to compute the q-characters of these modules.

3.2. **A closed system of type F_4.**

Theorem 3.4. *For $s \in \mathbb{Z}$ and $k, l, m, n \geq 1$, we have*

$$[\mathcal{T}^{(-2)}_{0,0,l-1,k}][\mathcal{T}^{(-2)}_{0,0,l,k}] = [\mathcal{T}^{(-2)}_{0,0,l,k-1}][\mathcal{T}^{(-2)}_{0,0,l-1,k+1}] + [\mathcal{T}^{(-2)}_{0,0,k+l,0}][\mathcal{T}^{(-2)}_{0,0,l-1,0}], \tag{3.1}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,0}][\widetilde{\mathcal{T}}^{(s)}_{n,m,0,0}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,0,0}][\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,0,0}] + [\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,m-1,0,0}][\widetilde{\mathcal{T}}^{(s)}_{0,n+m,0,0}], \tag{3.2}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,0}][\widetilde{\mathcal{T}}^{(s)}_{n,0,1,0}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,1,0}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,0}] + [\widetilde{\mathcal{T}}^{(s)}_{0,n,1,0}], \tag{3.3}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,l-2,0}][\widetilde{\mathcal{T}}^{(s)}_{n,0,l,0}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,l,0}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,l-2,0}] + [\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,0,l-2,0}][\widetilde{\mathcal{T}}^{(s)}_{0,n,l,0}],\ \ l \geq 2, \tag{3.4}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,0}][\widetilde{\mathcal{T}}^{(s)}_{0,m,1,0}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,1,0}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,0}] + [\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,0}][\widetilde{\mathcal{T}}^{(s)}_{0,0,1+2m,0}], \tag{3.5}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,l-2,0}][\widetilde{\mathcal{T}}^{(s)}_{0,m,l,0}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,l,0}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,l-2,0}] + [\widetilde{\mathcal{T}}^{(s+4)}_{m,0,l-2,0}][\widetilde{\mathcal{T}}^{(s)}_{0,0,l+2m,0}],\ \ l \geq 2, \tag{3.6}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,l,0}][\widetilde{\mathcal{T}}^{(s)}_{n,m,l,0}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,0}][\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,0}] + [\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,m-1,l,0}][\widetilde{\mathcal{T}}^{(s)}_{0,n+m,l,0}], \tag{3.7}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,0}][\widetilde{\mathcal{T}}^{(s)}_{n,0,0,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,0,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,0}] + [\widetilde{\mathcal{T}}^{(s)}_{0,n,0,k}],\ \ k = 1, 2, \tag{3.8}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,0}][\widetilde{\mathcal{T}}^{(s)}_{0,m,0,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,0,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,0}] + [\widetilde{\mathcal{T}}^{(s)}_{0,0,2m,k}][\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,0}],\ \ k = 1, 2, \tag{3.9}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-1}][\widetilde{\mathcal{T}}^{(s)}_{n,0,1,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,1,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,k-1}] + [\widetilde{\mathcal{T}}^{(s)}_{0,n,1,k}],\ \ k = 1, 2, \tag{3.10}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,l-2,k}][\widetilde{\mathcal{T}}^{(s)}_{n,0,l,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,l,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,l-2,k}] + [\widetilde{\mathcal{T}}^{(s)}_{0,n,l,k}][\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,0,l-2,k}],\ \ k = 1, 2,\ \ l \geq 2, \tag{3.11}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,k-1}][\widetilde{\mathcal{T}}^{(s)}_{0,m,1,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,1,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,k-1}] + [\widetilde{\mathcal{T}}^{(s)}_{0,0,1+2m,k}][\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,k-1}],\ \ k = 1, 2, \tag{3.12}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,l-2,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m,l,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,l,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,l-2,k}] + [\widetilde{\mathcal{T}}^{(s)}_{0,0,l+2m,k}][\widetilde{\mathcal{T}}^{(s+4)}_{m,0,l-2,k}],\ \ k = 1, 2,\ \ l \geq 2, \tag{3.13}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,k}][\widetilde{\mathcal{T}}^{(s)}_{n,m,0,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,0,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,0,k}] + [\widetilde{\mathcal{T}}^{(s)}_{0,n+m,0,k}][\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,0,k}],\ \ k = 1, 2, \tag{3.14}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,l,k}][\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,k}] + [\widetilde{\mathcal{T}}^{(s)}_{0,n+m,l,k}][\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,m-1,l,k}],\ \ k = 1, 2, \tag{3.15}$$

$$\begin{aligned}[\widetilde{\mathcal{P}}^{(s+2)}_{0,0,1,k-1}][\widetilde{\mathcal{P}}^{(s)}_{0,0,1,k}] = {} & [\widetilde{\mathcal{T}}^{(s+10)}_{0,0,0,k-4}][\widetilde{\mathcal{T}}^{(s+6)}_{0,0,0,k-2}][\widetilde{\mathcal{T}}^{(s+2)}_{0,0,0,k}][\widetilde{\mathcal{P}}^{(s)}_{0,0,2,k-1}] \\ & + [\widetilde{\mathcal{T}}^{(s+12)}_{0,0,0,k-5}][\widetilde{\mathcal{T}}^{(s+8)}_{0,0,0,k-3}][\widetilde{\mathcal{T}}^{(s+4)}_{0,0,0,k-1}][\widetilde{\mathcal{T}}^{(s)}_{0,0,0,k+1}][\widetilde{\mathcal{P}}^{(s+2)}_{0,1,0,k-2}],\ \ k \geq 3,\ \ l = 1,\end{aligned} \tag{3.16}$$

$$[\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l,k-1}][\widetilde{\mathcal{P}}^{(s)}_{0,0,l,k}] = [\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l-1,k}][\widetilde{\mathcal{P}}^{(s)}_{0,0,l+1,k-1}] + [\widetilde{\mathcal{T}}^{(s+2l+10)}_{0,0,0,k-5}][\widetilde{\mathcal{T}}^{(s)}_{0,0,0,k+l}][\widetilde{\mathcal{P}}^{(s+2)}_{0,\frac{l+1}{2},0,k-2}][\widetilde{\mathcal{P}}^{(s+4)}_{0,\frac{l-1}{2},0,k-1}], \tag{3.17}$$

where $k \geq 3$, l *is odd*, $l \geq 3$,

$$[\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l,k-1}][\widetilde{\mathcal{P}}^{(s)}_{0,0,l,k}] = [\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l-1,k}][\widetilde{\mathcal{P}}^{(s)}_{0,0,l+1,k-1}] + [\widetilde{\mathcal{T}}^{(s+2l+10)}_{0,0,0,k-5}][\widetilde{\mathcal{T}}^{(s)}_{0,0,0,k+l}][\widetilde{\mathcal{P}}^{(s+4)}_{0,\frac{l}{2},0,k-2}][\widetilde{\mathcal{P}}^{(s+2)}_{0,\frac{l}{2},0,k-1}], \tag{3.18}$$

where $k \geq 3$, l *is even*, $l \geq 2$,

$$[\widetilde{\mathcal{P}}^{(s+4)}_{0,1,0,k-2}][\widetilde{\mathcal{P}}^{(s)}_{0,1,0,k}] = [\widetilde{\mathcal{T}}^{(s+8)}_{0,0,0,k-2}][\widetilde{\mathcal{T}}^{(s+4)}_{0,0,0,k}][\widetilde{\mathcal{P}}^{(s)}_{0,2,0,k-2}] + [\widetilde{\mathcal{P}}^{(s)}_{0,0,2,k}][\widetilde{\mathcal{T}}^{(s+4)}_{1,0,0,k-2}], \ k \geq 3, \tag{3.19}$$

$$[\widetilde{\mathcal{P}}^{(s+4)}_{0,m,0,k-2}][\widetilde{\mathcal{P}}^{(s)}_{0,m,0,k}] = [\widetilde{\mathcal{P}}^{(s+4)}_{0,m-1,0,k}][\widetilde{\mathcal{P}}^{(s)}_{0,m+1,0,k-2}] + [\widetilde{\mathcal{P}}^{(s)}_{0,0,2m,k}][\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,k-2}], \ k \geq 3, \ m \geq 2, \tag{3.20}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-2}][\widetilde{\mathcal{T}}^{(s)}_{n,0,0,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,0,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,k-2}] + [\widetilde{\mathcal{P}}^{(s)}_{0,n,0,k}], \ k \geq 3. \tag{3.21}$$

Theorem 3.4 will be proved in Section 7. The system in Theorem 3.4 is a closed system in the sense that all modules in the system can be computed recursively using Kirillov-Reshetikhin modules.

Example 3.5. *The following are some equations in the system in Theorem 3.4.*

$[1_{-2}][1_{-4}2_{-1}] = [1_{-4}1_{-2}][2_{-1}] + [2_{-3}2_{-1}]$,
$[1_{-4}1_{-2}][1_{-6}1_{-4}2_{-1}] = [1_{-4}2_{-1}][1_{-6}1_{-4}1_{-2}] + [2_{-5}2_{-3}2_{-1}]$,
$[1_{-6}1_{-4}1_{-2}][1_{-8}1_{-6}1_{-4}2_{-1}] = [1_{-6}1_{-4}2_{-1}][1_{-8}1_{-6}1_{-4}1_{-2}] + [2_{-7}2_{-5}2_{-3}2_{-1}]$,
$[3_{-4}][1_{0}2_{-3}3_{-8}] = [1_{0}2_{-3}][3_{-8}3_{-4}] + [1_{0}2_{-7}2_{-5}2_{-3}][4_{-6}]$,
$[3_{-8}3_{-4}][1_{0}2_{-3}3_{-12}3_{-8} = [1_{0}2_{-3}3_{-8}][3_{-12}3_{-8}3_{-4}] + [1_{0}2_{-11}2_{-9}2_{-7}2_{-5}2_{-3}][4_{-10}4_{-6}]$,
$[3_{-12}3_{-8}3_{-4}][1_{0}2_{-3}3_{-16}3_{-12}3_{-8}] = [1_{0}2_{-3}3_{-12}3_{-8}][3_{-16}3_{-12}3_{-8}3_{-4}] + [1_{0}2_{-15}2_{-13}2_{-11}2_{-9}2_{-7}2_{-5}2_{-3}][4_{-14}4_{-10}4_{-6}]$,
$[4_{-6}][1_{0}2_{-3}4_{-10}] = [1_{0}2_{-3}][4_{-10}4_{-6}] + [1_{0}2_{-3}3_{-8}]$,
$[4_{-10}4_{-6}][1_{0}2_{-3}4_{-14}4_{-10}] = [1_{0}2_{-3}4_{-10}][4_{-14}4_{-10}4_{-6}] + [1_{0}2_{-3}3_{-12}3_{-8}]$,
$[4_{-14}4_{-10}4_{-6}][1_{0}2_{-3}4_{-18}4_{-14}4_{-10}] = [1_{0}2_{-3}4_{-14}4_{-10}][4_{-18}4_{-14}4_{-10}4_{-6}] + [1_{0}2_{-3}3_{-16}3_{-12}3_{-8}]$.

Moreover, we have the following theorem.

Theorem 3.6. *For each equation in Theorem* 3.4*, all summands on the right hand side are simple.*

Theorem 3.6 will be proved in Section 8.

3.3. **A system corresponding to the system in Theorem 3.4.**

Given $k, l, m, n \in \mathbb{Z}_{\geq 0}, s \in \mathbb{Z}$, let

$$\mathfrak{m}_{k,l,m,n} = \mathrm{Res}(\mathcal{T}^{(s)}_{k,l,m,n}) \text{ (resp. } \widetilde{\mathfrak{m}}_{k,l,m,n} = \mathrm{Res}(\widetilde{\mathcal{T}}^{(s)}_{k,l,m,n}))$$

be the restriction of $\mathcal{T}^{(s)}_{k,l,m,n}$ (resp. $\widetilde{\mathcal{T}}^{(s)}_{k,l,m,n}$) to $U_q\mathfrak{g}$. It is clear that $\mathrm{Res}(\mathcal{T}^{(s)}_{k,l,m,n})$ and $\mathrm{Res}(\widetilde{\mathcal{T}}^{(s)}_{k,l,m,n})$ do not depend on s. Let $\chi(M)$ (resp. $\chi(\widetilde{M})$) be the character of a $U_q\mathfrak{g}$-module M (resp. $\widetilde{M}$). By replacing each $[\mathcal{T}^{(s)}_{n,m,l,k}]$ (resp. $[\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}]$) in the system in Theorem 3.4 with $\chi(\mathfrak{m}_{n,m,l,k})$ (resp. $\chi(\widetilde{\mathfrak{m}}_{n,m,l,k})$), we obtain a system of equations consisting of the characters of $U_q\mathfrak{g}$-modules. The following are two equations in the system.

$$\chi(\mathfrak{m}_{0,0,l-1,k})\chi(\mathfrak{m}_{0,0,l,k}) = \chi(\mathfrak{m}_{0,0,l,k-1})\chi(\mathfrak{m}_{0,0,l-1,k+1}) + \chi(\mathfrak{m}_{0,0,k+l,0})\chi(\mathfrak{m}_{0,0,l-1,0}),$$
$$\chi(\widetilde{\mathfrak{m}}_{n,m-1,0,0})\chi(\widetilde{\mathfrak{m}}_{n,m,0,0}) = \chi(\widetilde{\mathfrak{m}}_{n-1,m,0,0})\chi(\widetilde{\mathfrak{m}}_{n+1,m-1,0,0}) + \chi(\widetilde{\mathfrak{m}}_{0,m-1,0,0})\chi(\widetilde{\mathfrak{m}}_{0,n+m,0,0}).$$

4. Relation between the system in Theorem 3.4 and cluster algebras

In this section, we will show that the equations in the system in Theorem 3.4 correspond to mutations in some cluster algebra $\mathscr{A}$. Moreover, every minimal affinization in the system in Theorem 3.4 corresponds to a cluster variable in the cluster algebra $\mathscr{A}$.

4.1. Definition of a cluster algebra $\mathscr{A}$.

Let $I = \{1, 2, 3, 4\}$ and

$$\begin{aligned} S &= \{-2u \mid u \in \mathbb{Z}_{\geq 0}\}, \\ S' &= \{-2u - 1 \mid u \in \mathbb{Z}_{\geq 0}\}. \end{aligned}$$

Let

$$V = (\{1\} \times S) \cup (\{2\} \times S') \cup (\{3\} \times S) \cup (\{4\} \times S).$$

We define Q with vertex set V as follows. The arrows of Q from the vertex (i, r) to the vertex (j, s) if and only if $b_{ij} \neq 0$ and $s = r - b_{ij} + d_i - d_j$. The quiver Q is the same as the quiver G^- of type F_4 defined in [HL13].

Let $\mathbf{t} = \mathbf{t}_1 \cup \mathbf{t}_2$, where

$$\mathbf{t}_1 = \{t^{(-2)}_{0,0,l,0},\ t^{(-4)}_{0,0,0,k} \mid k, l \in \mathbb{Z}_{\geq 1}\},$$

$$\mathbf{t}_2 = \{\widetilde{t}^{(-4n+4)}_{n,0,0,0},\ \widetilde{t}^{(-4m)}_{0,m,0,0},\ \widetilde{t}^{(-4n+2)}_{n,0,0,0},\ \widetilde{t}^{(-4m+2)}_{0,m,0,0},\ \widetilde{t}^{(-2l-2)}_{0,0,l,0},\ \widetilde{t}^{(2k-2)}_{0,0,0,k} \mid k, l, m, n \in \mathbb{Z}_{\geq 1}\}.$$

Let $\mathscr{A}$ be the cluster algebra defined by the initial seed $(\mathbf{t}, Q)$. By Definition 2.1, $\mathscr{A}$ is the $\mathbb{Q}$-subalgebra of the field of rational functions $\mathbb{Q}(\mathbf{t})$ generated by all the elements obtained from some elements of $\mathbf{t}$ via a finite sequence of seed mutations.

4.2. Mutation sequences.

We use "C_1", "C_2", "C_3", "C_4", "C_5", "C_6" to denote the column of vertices $(1, 0)$, $(1, 2)$, …, $(1, -2u)$, $\cdots$, the column of vertices $(2, -1)$, $(2, -3)$, …, $(2, -2u-1)$, $\cdots$, the column of vertices $(3, 0)$, $(3, -4)$, …, $(3, -4u)$, $\cdots$, the column of vertices $(3, -2)$, $(3, -6)$, …, $(3, -4u-2)$, $\cdots$, the column of vertices $(4, 0)$, $(4, -4)$, …, $(4, -4u)$, $\cdots$, the column of vertices $(4, -2)$, $(4, -6)$, …, $(4, -4u-2)$, $\cdots$, respectively in Q.

By saying that we mutate at the column C_i, $i \in \{1, 2, 3, 4, 5, 6\}$, we mean that we mutate the vertices of C_i as follows. First we mutate at the first vertex in this column, then the second vertex, and so on until the vertex at infinity. By saying that we mutate $(C_{i_0}, C_{i_1}, \ldots, C_{i_u})$, where $i_j \in \{1, 2, 3, 4, 5, 6\}$, $j = 0, 1, 2, \ldots, u$, we mean that we first mutate the column C_{i_1}, then the column C_{i_2}, and so on up to the column C_{i_u}.

For $k, l, m, n \in \mathbb{Z}_{\geq 1}$, we define some variables

$$\begin{gathered} t^{(-2)}_{0,0,l,k},\ \widetilde{t}^{(-4n-4m)}_{n,m,0,0},\ \widetilde{t}^{(-4n-4m+2)}_{n,m,0,0},\ \widetilde{t}^{(-4n-2l-2)}_{n,0,l,0},\ \widetilde{t}^{(-4m-2l-2)}_{0,m,l,0},\ \widetilde{t}^{(-4n-4m-2l-2)}_{n,m,l,0}, \\ \widetilde{t}^{(-4n-2l-2k-2)}_{n,0,l,k}\ (k \leq 2),\ \widetilde{t}^{(-4m-2l-2k-2)}_{0,m,l,k}\ (k \leq 2),\ \widetilde{t}^{(-4n-4m-2k-2)}_{n,m,0,k}\ (k \leq 2), \\ \widetilde{t}^{(-4n-4m-2l-2k-2)}_{n,m,l,k}\ (k \leq 2),\ \widetilde{p}^{(-2l-2k-2)}_{0,0,l,k},\ \widetilde{p}^{(-4m-2k-2)}_{0,m,0,k},\ \widetilde{t}^{(-4n-2k-2)}_{n,0,0,k}, \end{gathered} \tag{4.1}$$

recursively as follows. The variables in $\mathbf{t}$ are already defined.

We mutate the first vertex of the first C_1 in $(C_1, C_1, \ldots, C_1)$ from the initial seed, and define $t^{(-2)}_{0,0,0,1} = t'^{(-4)}_{0,0,0,1}$, we obtain a quiver (Q_{11}). Therefore

$$t^{(-2)}_{0,0,0,1} = t'^{(-4)}_{0,0,0,1} = \frac{t^{(-4)}_{0,0,0,2} + t^{(-2)}_{0,0,1,0}}{t^{(-4)}_{0,0,0,1}}. \tag{4.2}$$

We mutate the second vertex of the first C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,0,2} = t'^{(-4)}_{0,0,0,2}$, the quiver (Q_{11}) becomes a quiver (Q_{12}). Therefore

$$t^{(-2)}_{0,0,0,2} = t'^{(-4)}_{0,0,0,2} = \frac{t^{(-4)}_{0,0,0,3} t^{(-4)}_{0,0,0,1} + t^{(-2)}_{0,0,2,0}}{t^{(-4)}_{0,0,0,2}}. \tag{4.3}$$

We continue this procedure and mutate the vertices of the first C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,0,k} = t'^{(-4)}_{0,0,0,k}$ $(k = 3, 4, \ldots)$ recursively. Therefore

$$t^{(-2)}_{0,0,0,k} = t'^{(-4)}_{0,0,0,k} = \frac{t^{(-4)}_{0,0,0,k+1} t^{(-4)}_{0,0,0,k-1} + t^{(-2)}_{0,0,k,0}}{t^{(-4)}_{0,0,0,k}}, \quad k = 3, 4, \ldots \tag{4.4}$$

Now we finish the mutation of the first C_1 in $(C_1, C_1, \ldots, C_1)$.

We start to mutate the second C_1 in $(C_1, C_1, \ldots, C_1)$. We mutate the first vertex of the second C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,1,1} = t'^{(-2)}_{0,0,0,1}$, we obtain a quiver (Q_{21}). Therefore

$$t^{(-2)}_{0,0,1,1} = t'^{(-2)}_{0,0,0,1} = \frac{t^{(-2)}_{0,0,0,2} t^{(-2)}_{0,0,1,0} + t^{(-2)}_{0,0,2,0}}{t^{(-2)}_{0,0,0,1}}. \tag{4.5}$$

We mutate the second vertex of the second C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,1,2} = t'^{(-2)}_{0,0,0,2}$, the quiver (Q_{21}) becomes a quiver (Q_{22}). Therefore

$$t^{(-2)}_{0,0,1,2} = t'^{(-2)}_{0,0,0,2} = \frac{t^{(-2)}_{0,0,0,3} t^{(-2)}_{0,0,1,1} + t^{(-2)}_{0,0,3,0}}{t^{(-2)}_{0,0,0,2}}. \tag{4.6}$$

We continue this procedure and mutate the vertices of the second C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,1,k} = t'^{(-2)}_{0,0,0,k}$ $(k = 3, 4, \ldots)$ recursively. Therefore

$$t^{(-2)}_{0,0,1,k} = t'^{(-2)}_{0,0,0,k} = \frac{t^{(-2)}_{0,0,0,k+1} t^{(-2)}_{0,0,1,k-1} + t^{(-2)}_{0,0,k+1,0}}{t^{(-2)}_{0,0,0,k}}, \quad k = 3, 4, \ldots \tag{4.7}$$

Now we finish the mutation of the second C_1 in $(C_1, C_1, \ldots, C_1)$.

We start to mutate the third C_1 in $(C_1, C_1, \ldots, C_1)$. We mutate the first vertex of the third C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,2,1} = t'^{(-2)}_{0,0,1,1}$, we obtain a quiver (Q_{31}). Therefore

$$t^{(-2)}_{0,0,2,1} = t'^{(-2)}_{0,0,1,1} = \frac{t^{(-2)}_{0,0,1,2} t^{(-2)}_{0,0,2,0} + t^{(-2)}_{0,0,1,0} t^{(-2)}_{0,0,3,0}}{t^{(-2)}_{0,0,1,1}}. \tag{4.8}$$

We mutate the second vertex of the third C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,2,2} = t'^{(-2)}_{0,0,1,2}$, the quiver (Q_{31}) becomes a quiver (Q_{32}). Therefore

$$t^{(-2)}_{0,0,2,2} = t'^{(-2)}_{0,0,1,2} = \frac{t^{(-2)}_{0,0,1,3} t^{(-2)}_{0,0,2,1} + t^{(-2)}_{0,0,1,0} t^{(-2)}_{0,0,4,0}}{t^{(-2)}_{0,0,1,2}}. \tag{4.9}$$

We continue this procedure and mutate vertices of the third C_1 in $(C_1, C_1, \ldots, C_1)$ and define $t^{(-2)}_{0,0,2,k} = t'^{(-2)}_{0,0,1,k}$ $(k = 3, 4, \ldots)$ recursively. Therefore

$$t^{(-2)}_{0,0,2,k} = t'^{(-2)}_{0,0,1,k} = \frac{t^{(-2)}_{0,0,1,k+1} t^{(-2)}_{0,0,2,k-1} + t^{(-2)}_{0,0,1,0} t^{(-2)}_{0,0,k+2,0}}{t^{(-2)}_{0,0,1,k}}, \quad k = 3, 4, \ldots \tag{4.10}$$

Now we finish the mutation of the third C_1 in $(C_1, C_1, \ldots, C_1)$. We continue this procedure and mutate the $(d+1)$-th C_1 $(d = 3, 5, \ldots, l)$ in order. We define $t^{(-2)}_{0,0,d,k} = t'^{(-2)}_{0,0,d-1,k}$, where $(0,0,d,k)$=$\{(0,0,3,1), (0,0,3,2), (0,0,3,3), (0,0,3,4), \ldots; (0,0,4,1), (0,0,4,2), (0,0,4,3), (0,0,4,4) \ldots; (0,0,5,1), (0,0,5,2), (0,0,5,3), (0,0,5,4), \ldots; (0,0,l,1), (0,0,l,2), (0,0,l,3), (0,0,l, 4), \ldots\}$ recursively. Therefore

$$t^{(-2)}_{0,0,d,k} = t'^{(-2)}_{0,0,d-1,k} = \frac{t^{(-2)}_{0,0,d-1,k+1} t^{(-2)}_{0,0,d,k-1} + t^{(-2)}_{0,0,d-1,0} t^{(-2)}_{0,0,k+d,0}}{t^{(-2)}_{0,0,d-1,k}}. \tag{4.11}$$

We write the definition of the variables in (4.1) and the corresponding mutation sequences in Table 1.

In Table 1, we use

$$\underbrace{C_{i_1}, C_{i_2}, \cdots, C_{i_u}, C_{i_1}, C_{i_2}, \cdots, C_{i_u}, \cdots, C_{i_1}, C_{i_2}, \cdots, C_{i_u}}_{m},$$

where $m \in \mathbb{Z}_{\geq 1}$, $u \in \mathbb{Z}_{\geq 1}$, to denote the mutation sequences

$$C_{i_1}, C_{i_2}, \cdots, C_{i_u}, C_{i_1}, C_{i_2}, \cdots, C_{i_u}, \cdots, C_{i_1}, C_{i_2}, \cdots, C_{i_u},$$

where the number of C_{i_j} $(j \in \{1, 2, \cdots, u\})$ is m.

Mutation sequences	Definition of variables in (4.1) and mutation equations
$(\underbrace{C_1, C_1, \ldots, C_1}_{l+1})$	$t^{(-2)}_{0,0,l,k} = t'^{(-2)}_{0,0,l-1,k} = \frac{t^{(-2)}_{0,0,l-1,k+1} t^{(-2)}_{0,0,l,k-1} + t^{(-2)}_{0,0,l-1,0} t^{(-2)}_{0,0,k+l,0}}{t^{(-2)}_{0,0,l-1,k}} \quad (1)$
$(\underbrace{C_5, C_5, \ldots, C_5}_{m+1})$ $(\underbrace{C_6, C_6, \ldots, C_6}_{m})$	$\widetilde{t}^{(-4n-4m)}_{n,m,0,0} = \widetilde{t}'^{(-4n-4m+4)}_{n,m-1,0,0} = \frac{\widetilde{t}^{(-4n-4m)}_{n+1,m-1,0,0} \widetilde{t}^{(-4n-4m+4)}_{n-1,m,0,0} + \widetilde{t}^{(-4m+4)}_{0,m-1,0,0} \widetilde{t}^{(-4n-4m)}_{0,n+m,0,0}}{\widetilde{t}^{(-4n-4m+4)}_{n,m-1,0,0}} \quad (2)$ $\widetilde{t}^{(-4n-4m+2)}_{n,m,0,0} = \widetilde{t}'^{(-4n-4m+6)}_{n,m-1,0,0} = \frac{\widetilde{t}^{(-4n-4m+2)}_{n+1,m-1,0,0} \widetilde{t}^{(-4n-4m+6)}_{n-1,m,0,0} + \widetilde{t}^{(-4m+6)}_{0,m-1,0,0} \widetilde{t}^{(-4n-4m+2)}_{0,n+m,0,0}}{\widetilde{t}^{(-4n-4m+6)}_{n,m-1,0,0}} \quad (3)$
$(C_5,\underbrace{C_4, C_5, \ldots, C_4, C_5}_{\frac{l+1}{2},\ l \text{ is odd}})$ $(\underbrace{C_3, C_6, \ldots, C_3, C_6}_{\frac{l+2}{2},\ l \text{ is even}})$	$\widetilde{t}^{(-4n-4)}_{n,0,1,0} = \widetilde{t}'^{(-4n)}_{n,0,0,0} = \frac{\widetilde{t}^{(-4n-4)}_{n+1,0,0,0} \widetilde{t}^{(-4n)}_{n-1,0,1,0} + \widetilde{t}^{(-4n-4)}_{0,n,1,0}}{\widetilde{t}^{(-4n)}_{n,0,0,0}} \quad (4)$ $\widetilde{t}^{(-4n-2l-2)}_{n,0,l,0} = \widetilde{t}'^{(-4n-2l+2)}_{n,0,l-2,0} = \frac{\widetilde{t}^{(-4n-2l-2)}_{n+1,0,l-2,0} \widetilde{t}^{(-4n-2l+2)}_{n-1,0,l,0} + \widetilde{t}^{(-4n-2l-2)}_{0,n,l,0} \widetilde{t}^{(-2l+2)}_{0,0,l-2,0}}{\widetilde{t}^{(-4l-2l+2)}_{n,0,l-2,0}} \quad (5)$ $\widetilde{t}^{(-4m-4)}_{0,m,1,0} = \widetilde{t}'^{(-4m)}_{0,m,0,0} = \frac{\widetilde{t}^{(-4m-4)}_{0,m+1,0,0} \widetilde{t}^{(-4m)}_{0,m-1,1,0} + \widetilde{t}^{(-4m-4)}_{0,0,2m+1,0} \widetilde{t}^{(-4m)}_{m,0,0,0}}{\widetilde{t}^{(-4m)}_{0,m,0,0}} \quad (6)$ $\widetilde{t}^{(-4m-2l-2)}_{0,m,l,0} = \widetilde{t}'^{(-4m-2l+2)}_{0,m,l-2,0} = \frac{\widetilde{t}^{(-4m-2l-2)}_{0,m+1,l-2,0} \widetilde{t}^{(-4m-2l+2)}_{0,m-1,l,0} + \widetilde{t}^{(-4m-2l-2)}_{0,0,2m+l,0} \widetilde{t}^{(-4m-2l+2)}_{m,0,l-2,0}}{\widetilde{t}^{(-4m-2l+2)}_{0,m,l-2,0}} \quad (7)$
$(C_5,\underbrace{C_4, C_5, \ldots, C_4, C_5}_{\frac{l+1}{2},\ l \text{ is odd}},\underbrace{C_5, \ldots, C_5}_{m})$ $(\underbrace{C_3, C_6, \ldots, C_3, C_6}_{\frac{l+2}{2},\ l \text{ is even}},\underbrace{C_6, \ldots, C_6}_{m})$	$\widetilde{t}^{(-4n-4m-2l-2)}_{n,m,l,0} = \widetilde{t}'^{(-4n-4m-2l+2)}_{n,m-1,l,0} = \frac{\widetilde{t}^{(-4n-4m-2l-2)}_{n+1,m-1,l,0} \widetilde{t}^{(-4n-4m-2l+2)}_{n-1,m,l,0} + \widetilde{t}^{(-4n-4m-2l-2)}_{0,n+m,l,0} \widetilde{t}^{(-4m-2l+2)}_{0,m-1,l,0}}{\widetilde{t}^{(-4n-4m-2l+2)}_{n,m-1,l,0}} \quad (8)$
$(C_3,C_6,C_2,C_5,C_4,C_5)$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_3,C_6)$	$\widetilde{t}^{(-4n-2k-2)}_{n,0,0,k} = \widetilde{t}'^{(-4n-2k+2)}_{n,0,0,0} = \frac{\widetilde{t}^{(-4n-2k+2)}_{n-1,0,0,k} \widetilde{t}^{(-4n-2k-2)}_{n+1,0,0,0} + \widetilde{t}^{(-4n-2k-2)}_{0,n,0,k}}{\widetilde{t}^{(-4n-2k+2)}_{n,0,0,0}}, \quad (9)$ $\widetilde{t}^{(-4m-2k-2)}_{0,m,0,k} = \widetilde{t}'^{(-4m-2k+2)}_{0,m,0,0} = \frac{\widetilde{t}^{(-4m-2k+2)}_{0,m-1,0,k} \widetilde{t}^{(-4m-2k-2)}_{0,m+1,0,0} + \widetilde{t}^{(-4m-2k-2)}_{0,0,2m,k} \widetilde{t}^{(-4m-2k+2)}_{m,0,0,0}}{\widetilde{t}^{(-4m-2k+2)}_{0,m,0,0}},\ k{=}1,2, \quad (10)$
$(C_3,C_6,C_2,\underbrace{C_3, C_6, \ldots, C_3, C_6}_{\frac{l+1}{2},\ l \text{ is odd}})$ $(C_3,C_6,C_2,C_5,C_4,C_5,\underbrace{C_4, C_5, \ldots, C_4, C_5}_{\frac{l+2}{2},\ l \text{ is even}})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,\underbrace{C_4, C_5, \ldots, C_4, C_5}_{\frac{l+1}{2},\ l \text{ is odd}})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_3,C_6,\underbrace{C_3, C_6, \ldots, C_3, C_6}_{\frac{l}{2},\ l \text{ is even}})$	$\widetilde{t}^{(-4n-2k-4)}_{n,0,1,k} = \widetilde{t}'^{(-4n-2k)}_{n,0,0,k-1} = \frac{\widetilde{t}^{(-4n-2k)}_{n-1,0,1,k} \widetilde{t}^{(-4n-2k-4)}_{n+1,0,0,k-1} + \widetilde{t}^{(-4n-2k-4)}_{0,n,1,k} \widetilde{t}^{(-2k)}_{0,0,0,k-1}}{\widetilde{t}^{(-4n-2n)}_{n,0,0,k-1}},\ k{=}1,2,\ l{=}1; \quad (11)$ $\widetilde{t}^{(-4n-2l-2k-2)}_{n,0,l,k} = \widetilde{t}'^{(-4n-2l-2k+2)}_{n,0,l-2,k} = \frac{\widetilde{t}^{(-4n-2l-2k+2)}_{n-1,0,l,k} \widetilde{t}^{(-4n-2l-2k-2)}_{n+1,0,l-2,k} + \widetilde{t}^{(-4n-2l-2k-2)}_{0,n,l,k} \widetilde{t}^{(-2l-2k+2)}_{0,0,l-2,k}}{\widetilde{t}^{(-4n-2l-2n+2)}_{n,0,l-2,k}},\ k{=}1,2,\ l{\geq}2, \quad (12)$ $\widetilde{t}^{(-4m-2k-4)}_{0,m,1,k} = \widetilde{t}'^{(-4m-2k)}_{0,m,0,k-1} = \frac{\widetilde{t}^{(-4m-2k)}_{0,m-1,1,k} \widetilde{t}^{(-4m-2k-4)}_{0,m+1,0,k-1} + \widetilde{t}^{(-4m-2k-4)}_{0,0,2m+1,k} \widetilde{t}^{(-4m-2k)}_{m,0,0,k-1}}{\widetilde{t}^{(-4m-2k)}_{0,m,0,k-1}},\ k{=}1,2,\ l{=}1, \quad (13)$ $\widetilde{t}^{(-4m-2l-2k-2)}_{0,m,l,k} = \widetilde{t}'^{(-4m-2l-2k+2)}_{0,m,l-2,k} = \frac{\widetilde{t}^{(-4m-2l-2k+2)}_{0,m-1,l,k} \widetilde{t}^{(-4m-2l-2k-2)}_{0,m+1,l-2,k} + \widetilde{t}^{(-4m-2l-2k-2)}_{0,0,2m+l,k} \widetilde{t}^{(-4m-2l-2k+2)}_{m,0,l-2,k}}{\widetilde{t}^{(-4m-2l-2k+2)}_{0,m,l-2,k}},\ k{=}1,2,\ l{\geq}2, \quad (14)$
$(C_3,C_6,C_2,C_5,C_4,C_5,\underbrace{C_5, \ldots, C_5}_{\frac{m+1}{2},\ m \text{ is odd}})$ $(C_3,C_6,C_2,C_3,C_6,\underbrace{C_6, \ldots, C_6}_{\frac{m}{2},\ m \text{ is even}})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_3,C_6,\underbrace{C_6, \ldots, C_6}_{\frac{m+1}{2},\ m \text{ is odd}})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_4,C_5,\underbrace{C_5, \ldots, C_5}_{\frac{m}{2},\ m \text{ is even}})$	$\widetilde{t}^{(-4n-4m-2k-2)}_{n,m,0,k} = \widetilde{t}'^{(-4n-4m-2k+2)}_{n,m-1,0,k} = \frac{\widetilde{t}^{(-4n-4m-2k+2)}_{n-1,m,0,k} \widetilde{t}^{(-4n-4m-2k-2)}_{n+1,m-1,0,k} + \widetilde{t}^{(-4n-4m-2k-2)}_{0,n+m,0,k} \widetilde{t}^{(-4m-2k+2)}_{0,m-1,0,k}}{\widetilde{t}^{(-4n-4m-2k+2)}_{n,m-1,0,k}},\ k{=}1,2, \quad (15)$
$(C_3,C_6,C_2,\underbrace{C_3, C_6, \ldots, C_3, C_6}_{\frac{l+1}{2},\ l \text{ is odd}},\underbrace{C_6, \ldots, C_6}_{m})$ $(C_3,C_6,C_2,C_5,C_4,C_5,\underbrace{C_4, C_5, \ldots, C_4, C_5}_{\frac{l}{2},\ l \text{ is even}},\underbrace{C_5, \ldots, C_5}_{m})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_4,C_5,\underbrace{C_4, C_5, \ldots, C_4, C_5}_{\frac{l+1}{2},\ l \text{ is odd}},\underbrace{C_5, \ldots, C_5}_{m})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_3,C_6,\underbrace{C_3, C_6, \ldots, C_3, C_6}_{\frac{l}{2},\ l \text{ is even}},\underbrace{C_6, \ldots, C_6}_{m})$	$\widetilde{t}^{(-4n-4m-2l-2k-2)}_{n,m,l,k} = \widetilde{t}'^{(-4n-4m-2l-2k+2)}_{n,m-1,l,k} = \frac{\widetilde{t}^{(-4n-4m-2l-2k-2)}_{n+1,m-1,l,k} \widetilde{t}^{(-4n-4m-2l-2k+2)}_{n-1,m,l,k} + \widetilde{t}^{(-4n-4m-2l-2k-2)}_{0,n+m,l,k} \widetilde{t}^{(-4m-2l-2k+2)}_{0,m-1,l,k}}{\widetilde{t}^{(-4n-4m-2l-2k+2)}_{n,m-1,l,k}},\ k{=}1,2, \quad (16)$
$(C_3,C_6,C_2,C_5,C_4,C_5,C_2,\underbrace{C_3, C_6, C_2, C_4, C_5, C_2, \cdots, C_3, C_6, C_2, C_4, C_5, C_2}_{k-2})$	$\widetilde{p}^{(-2k-4)}_{0,0,1,k} = \widetilde{p}'^{(-2k-2)}_{0,0,1,k-1} = \frac{\widetilde{t}^{(-2k+6)}_{0,0,0,k-4} \widetilde{t}^{(-2k+2)}_{0,0,0,k-2} \widetilde{t}^{(-2k-2)}_{0,0,0,k} \widetilde{p}^{(-2k-4)}_{0,0,2,k-1} + \widetilde{t}^{(-2k+8)}_{0,0,0,k-5} \widetilde{t}^{(-2k+4)}_{0,0,0,k-3} \widetilde{t}^{(-2k)}_{0,0,0,k-1} \widetilde{t}^{(-2k-4)}_{0,0,0,k+1} \widetilde{p}^{(-2k-2)}_{0,1,0,k-2}}{\widetilde{p}^{(-2k-2)}_{0,0,1,k-1}},\ k{\geq}3,\ l{=}1, \quad (17)$ $\widetilde{p}^{(-2k-2l-2)}_{0,0,l,k} = \widetilde{p}'^{(-2k-2l)}_{0,0,l,k-1} = \frac{\widetilde{p}^{(-2k-2l)}_{0,0,l-1,k} \widetilde{p}^{(-2k-2l-2)}_{0,0,l+1,k-1} + \widetilde{t}^{(-2k+8)}_{0,0,0,k-5} \widetilde{t}^{(-2k-2l-2)}_{0,0,0,k+l} \widetilde{p}^{(-2k-2l)}_{0,\frac{l+1}{2},0,k-2} \widetilde{p}^{(-2k-2l+2)}_{0,\frac{l-1}{2},0,k-1}}{\widetilde{p}^{(-2k-2l)}_{0,0,l,k-1}},\ k{\geq}3,\ l \text{ is odd},\ l{\geq}3, \quad (18)$ $\widetilde{p}^{(-2k-2l-2)}_{0,0,l,k} = \widetilde{p}'^{(-2k-2l)}_{0,0,l,k-1} = \frac{\widetilde{p}^{(-2k-2l)}_{0,0,l-1,k} \widetilde{p}^{(-2k-2l-2)}_{0,0,l+1,k-1} + \widetilde{t}^{(-2k+8)}_{0,0,0,k-5} \widetilde{t}^{(-2k-2l-2)}_{0,0,0,k+l} \widetilde{p}^{(-2k-2l+2)}_{0,\frac{l}{2},0,k-2} \widetilde{p}^{(-2k-2l)}_{0,\frac{l}{2},0,k-1}}{\widetilde{p}^{(-2k-2l)}_{0,0,l,k-1}},\ k{\geq}3,\ l \text{ is even},\ l{\geq}2, \quad (19)$
$(C_3,C_6,C_2,C_5,C_4,C_5,\underbrace{C_2, C_3, C_6, C_2, C_4, C_5, \cdots, C_2, C_3, C_6, C_2, C_4, C_5}_{\frac{k-1}{2},\ k \text{ is odd}})$ $(C_3,C_6,C_2,C_5,C_4,C_5,C_2,C_3,C_6,\underbrace{C_2, C_4, C_5, C_2, C_3, C_6, \cdots, C_2, C_4, C_5, C_2, C_3, C_6}_{\frac{k-2}{2},\ k \text{ is even}})$	$\widetilde{p}^{(-2k-6)}_{0,1,0,k} = \widetilde{p}'^{(-2k-2)}_{0,1,0,k-2} = \frac{\widetilde{t}^{(-2k+2)}_{0,0,0,k-2} \widetilde{t}^{(s+4)}_{0,0,0,k} \widetilde{p}^{(-2k-6)}_{0,2,0,k-2} + \widetilde{p}^{(-2k-6)}_{0,0,2,k} \widetilde{t}^{(-2k-2)}_{1,0,0,k-2}}{\widetilde{p}^{(-2k-2)}_{0,1,0,k-2}},\ k{\geq}3,\ m{=}1, \quad (20)$ $\widetilde{p}^{(-4m-2k-2)}_{0,m,0,k} = \widetilde{p}'^{(-4m-2k+2)}_{0,m,0,k-2} = \frac{\widetilde{p}^{(-4m-2k+2)}_{0,m-1,0,k} \widetilde{p}^{(-4m-2k-2)}_{0,m+1,0,k-2} + \widetilde{p}^{(-4m-2k-2)}_{0,0,2m,k} \widetilde{t}^{(-4m-2k+2)}_{m,0,0,k-2}}{\widetilde{p}^{(-4m-2k+2)}_{0,m,0,k-2}},\ k{\geq}3,\ m{\geq}2, \quad (21)$ $\widetilde{t}^{(-4n-2k-2)}_{n,0,0,k} = \widetilde{t}'^{(-4n-2k+2)}_{n,0,0,k-2} = \frac{\widetilde{t}^{(-4n-2k+2)}_{n-1,0,0,k} \widetilde{t}^{(-4n-2k-2)}_{n+1,0,0,k-2} + \widetilde{p}^{(-4n-2k-2)}_{0,n,0,k}}{\widetilde{t}^{(-2k-2n+2)}_{n,0,0,k-2}},\ k{\geq}3. \quad (22)$

TABLE 1. Mutation sequences.

4.3. **The equations in the system of type F_4 correspond to mutations in the cluster algebra $\mathscr{A}$.** Equation (1) corresponds to Equation (3.1) in Theorem 3.4, Equations (2) and (3) correspond to Equation (3.2) in Theorem 3.4. Equations (4)–(22) correspond to Equations (3.3)–(3.21) in Theorem 3.4 respectively. Therefore we have the following theorem.

Theorem 4.1. *Each minimal affinizations in Theorem 3.4 corresponds to a cluster variable in $\mathscr{A}$ defined in Section 4.1.*

5. THE DUAL SYSTEM OF THEOREM 3.4

In this section, we study the dual system of Theorem 3.4.

Theorem 5.1 (Theorem 3.9, [Her07]). *For $l, m, n \in \mathbb{Z}_{\geq 1}$, $s \in \mathbb{Z}$, the modules $\mathcal{T}^{(s)}_{n,m,0,0}$, $\mathcal{T}^{(s)}_{n,0,l,0}$, $\mathcal{T}^{(s)}_{0,m,l,0}$, $\mathcal{T}^{(s)}_{n,m,l,0}$ are anti-special.*

We have the following theorem.

Theorem 5.2. *The modules*

$$\widetilde{\mathcal{T}}^{(s)}_{0,0,l,k},\ \mathcal{T}^{(s)}_{n,0,l,0},\ \mathcal{T}^{(s)}_{0,m,l,0},\ \mathcal{T}^{(s)}_{n,m,0,0},\ \mathcal{T}^{(s)}_{n,m,l,0},\ \mathcal{T}^{(s)}_{n,0,0,k}, \mathcal{T}^{(s)}_{n,m,0,k}(k \leq 2),$$
$$\mathcal{T}^{(s)}_{n,m,l,k}(k \leq 2),\ \mathcal{T}^{(s)}_{n,0,l,k}(k \leq 2),\ \mathcal{T}^{(s)}_{0,m,l,k}(k \leq 2),\ \mathcal{P}^{(s)}_{0,0,l,k},\ \mathcal{P}^{(s)}_{0,m,0,k},$$

where $k, l, m, n \in \mathbb{Z}_{\geq 1}$, $s \in \mathbb{Z}$, are anti-special.

Proof. The proof of the theorem follows from dual arguments in the proof of Theorem 3.3. □

Lemma 5.3. *Let $\iota : \mathbb{Z}\mathcal{P} \to \mathbb{Z}\mathcal{P}$ be a homomorphism of rings such that $Y_{1,aq^s} \mapsto Y^{-1}_{1,aq^{18-s}}$, $Y_{2,aq^s} \mapsto Y^{-1}_{2,aq^{18-s}}$, $Y_{3,aq^s} \mapsto Y^{-1}_{3,aq^{18-s}}$, $Y_{4,aq^s} \mapsto Y^{-1}_{4,aq^{18-s}}$ for all $a \in \mathbb{C}^\times, s \in \mathbb{Z}$. Then*

$$\chi_q(\widetilde{\mathcal{T}}^{(s)}_{k,l,m,n}) = \iota(\chi_q(\mathcal{T}^{(s)}_{k,l,m,n})), \quad \chi_q(\mathcal{T}^{(s)}_{k,l,m,n}) = \iota(\chi_q(\widetilde{\mathcal{T}}^{(s)}_{k,l,m,n})).$$

Proof. The proof is similar to Lemma 7.3 in [LM13]. □

Theorem 5.4. *For $s \in \mathbb{Z}$, $k, l, m, n \in \mathbb{Z}_{\geq 0}$, we have the following system of equations.*

$$[\widetilde{\mathcal{T}}^{(-2)}_{0,0,l-1,k}][\widetilde{\mathcal{T}}^{(-2)}_{0,0,l,k}] = [\widetilde{\mathcal{T}}^{(-2)}_{0,0,l,k-1}][\widetilde{\mathcal{T}}^{(-2)}_{0,0,l-1,k+1}] + [\widetilde{\mathcal{T}}^{(-2)}_{0,0,k+l,0}][\widetilde{\mathcal{T}}^{(-2)}_{0,0,l-1,0}], \tag{5.1}$$

$$[\mathcal{T}^{(s+4)}_{n,m-1,0,0}][\mathcal{T}^{(s)}_{n,m,0,0}] = [\mathcal{T}^{(s+4)}_{n-1,m,0,0}][\mathcal{T}^{(s)}_{n+1,m-1,0,0}] + [\mathcal{T}^{(s+4n+4)}_{0,m-1,0,0}][\mathcal{T}^{(s)}_{0,n+m,0,0}], \tag{5.2}$$

$$[\mathcal{T}^{(s+4)}_{n,0,0,0}][\mathcal{T}^{(s)}_{n,0,1,0}] = [\mathcal{T}^{(s+4)}_{n-1,0,1,0}][\mathcal{T}^{(s)}_{n+1,0,0,0}] + [\mathcal{T}^{(s)}_{0,n,1,0}], \tag{5.3}$$

$$[\mathcal{T}^{(s+4)}_{n,0,l-2,0}][\mathcal{T}^{(s)}_{n,0,l,0}] = [\mathcal{T}^{(s+4)}_{n-1,0,l,0}][\mathcal{T}^{(s)}_{n+1,0,l-2,0}] + [\mathcal{T}^{(s+4n+4)}_{0,0,l-2,0}][\mathcal{T}^{(s)}_{0,n,l,0}],\ l \geq 2, \tag{5.4}$$

$$[\mathcal{T}^{(s+4)}_{0,m,0,0}][\mathcal{T}^{(s)}_{0,m,1,0}] = [\mathcal{T}^{(s+4)}_{0,m-1,1,0}][\mathcal{T}^{(s)}_{0,m+1,0,0}] + [\mathcal{T}^{(s+4)}_{m,0,0,0}][\mathcal{T}^{(s)}_{0,0,1+2m,0}], \tag{5.5}$$

$$[\mathcal{T}^{(s+4)}_{0,m,l-2,0}][\mathcal{T}^{(s)}_{0,m,l,0}] = [\mathcal{T}^{(s+4)}_{0,m-1,l,0}][\mathcal{T}^{(s)}_{0,m+1,l-2,0}] + [\mathcal{T}^{(s+4)}_{m,0,l-2,0}][\mathcal{T}^{(s)}_{0,0,l+2m,0}],\ l \geq 2 \tag{5.6}$$

$$[\mathcal{T}^{(s+4)}_{n,m-1,l,0}][\mathcal{T}^{(s)}_{n,m,l,0}] = [\mathcal{T}^{(s+4)}_{n-1,m,l,0}][\mathcal{T}^{(s)}_{n+1,m-1,l,0}] + [\mathcal{T}^{(s+4n+4)}_{0,m-1,l,0}][\mathcal{T}^{(s)}_{0,n+m,l,0}], \tag{5.7}$$

$$[\mathcal{T}^{(s+4)}_{n,0,0,0}][\mathcal{T}^{(s)}_{n,0,0,k}] = [\mathcal{T}^{(s+4)}_{n-1,0,0,k}][\mathcal{T}^{(s)}_{n+1,0,0,0}] + [\mathcal{T}^{(s)}_{0,n,0,k}],\ k = 1, 2, \tag{5.8}$$

$$[\mathcal{T}^{(s+4)}_{0,m,0,0}][\mathcal{T}^{(s)}_{0,m,0,k}] = [\mathcal{T}^{(s+4)}_{0,m-1,0,k}][\mathcal{T}^{(s)}_{0,m+1,0,0}] + [\mathcal{T}^{(s)}_{0,0,2m,k}][\mathcal{T}^{(s+4)}_{m,0,0,0}], \ k = 1, 2, \tag{5.9}$$

$$[\mathcal{T}^{(s+4)}_{n,0,0,k-1}][\mathcal{T}^{(s)}_{n,0,1,k}] = [\mathcal{T}^{(s+4)}_{n-1,0,1,k}][\mathcal{T}^{(s)}_{n+1,0,0,k-1}] + [\mathcal{T}^{(s)}_{0,n,1,k}], \ k = 1, 2, \tag{5.10}$$

$$[\mathcal{T}^{(s+4)}_{0,m,0,k-1}][\mathcal{T}^{(s)}_{0,m,1,k}] = [\mathcal{T}^{(s+4)}_{0,m-1,1,k}][\mathcal{T}^{(s)}_{0,m+1,0,k-1}] + [\mathcal{T}^{(s)}_{0,0,1+2m,k}][\mathcal{T}^{(s+4)}_{m,0,0,k-1}], \ k = 1, 2, \tag{5.11}$$

$$[\mathcal{T}^{(s+4)}_{n,0,l-2,k}][\mathcal{T}^{(s)}_{n,0,l,k}] = [\mathcal{T}^{(s+4)}_{n-1,0,l,k}][\mathcal{T}^{(s)}_{n+1,0,l-2,k}] + [\mathcal{T}^{(s)}_{0,n,l,k}][\mathcal{T}^{(s+4n+4)}_{0,0,l-2,k}], \ k = 1, 2, \ l \geq 2, \tag{5.12}$$

$$[\mathcal{T}^{(s+4)}_{0,m,l-2,k}][\mathcal{T}^{(s)}_{0,m,l,k}] = [\mathcal{T}^{(s+4)}_{0,m-1,l,k}][\mathcal{T}^{(s)}_{0,m+1,l-2,k}] + [\mathcal{T}^{(s)}_{0,0,l+2m,k}][\mathcal{T}^{(s+4)}_{m,0,l-2,k}], \ k = 1, 2, \ l \geq 2, \tag{5.13}$$

$$[\mathcal{T}^{(s+4)}_{n,m-1,0,k}][\mathcal{T}^{(s)}_{n,m,0,k}] = [\mathcal{T}^{(s+4)}_{n-1,m,0,k}][\mathcal{T}^{(s)}_{n+1,m-1,0,k}] + [\mathcal{T}^{(s)}_{0,n+m,0,k}][\mathcal{T}^{(s+4)}_{0,m-1,0,k}], \ k = 1, 2, \tag{5.14}$$

$$[\mathcal{T}^{(s+4)}_{n,m-1,l,k}][\mathcal{T}^{(s)}_{n,m,l,k}] = [\mathcal{T}^{(s+4)}_{n-1,m,l,k}][\mathcal{T}^{(s)}_{n+1,m-1,l,k}] + [\mathcal{T}^{(s)}_{0,n+m,l,k}][\mathcal{T}^{(s+4n+4)}_{0,m-1,l,k}], \ k = 1, 2, \tag{5.15}$$

$$\begin{aligned} [\mathcal{P}^{(s+2)}_{0,0,1,k-1}][\mathcal{P}^{(s)}_{0,0,1,k}] &= [\mathcal{T}^{(s+10)}_{0,0,0,k-4}][\mathcal{T}^{(s+6)}_{0,0,0,k-2}][\mathcal{T}^{(s+2)}_{0,0,0,k}][\mathcal{P}^{(s)}_{0,0,2,k-1}] \\ &\quad + [\mathcal{T}^{(s+12)}_{0,0,0,k-5}][\mathcal{T}^{(s+8)}_{0,0,0,k-3}][\mathcal{T}^{(s+4)}_{0,0,0,k-1}][\mathcal{T}^{(s)}_{0,0,0,k+1}][\mathcal{P}^{(s+2)}_{0,1,0,k-2}], \ k \geq 3, \end{aligned} \tag{5.16}$$

$$[\mathcal{P}^{(s+2)}_{0,0,l,k-1}][\mathcal{P}^{(s)}_{0,0,l,k}] = [\mathcal{P}^{(s+2)}_{0,0,l-1,k}][\mathcal{P}^{(s)}_{0,0,l+1,k-1}] + [\mathcal{T}^{(s+2l+10)}_{0,0,0,k-5}][\mathcal{T}^{(s)}_{0,0,0,k+l}][\mathcal{P}^{(s+2)}_{0,\frac{l+1}{2},0,k-2}][\mathcal{P}^{(s+4)}_{0,\frac{l-1}{2},0,k-1}], \tag{5.17}$$

where $k \geq 3$, l *is odd,* $l \geq 3$,

$$[\mathcal{P}^{(s+2)}_{0,0,l,k-1}][\mathcal{P}^{(s)}_{0,0,l,k}] = [\mathcal{P}^{(s+2)}_{0,0,l-1,k}][\mathcal{P}^{(s)}_{0,0,l+1,k-1}] + [\mathcal{T}^{(s+2l+10)}_{0,0,0,k-5}][\mathcal{T}^{(s)}_{0,0,0,k+l}][\mathcal{P}^{(s+4)}_{0,\frac{l}{2},0,k-2}][\mathcal{P}^{(s+2)}_{0,\frac{l}{2},0,k-1}], \tag{5.18}$$

where $k \geq 3$, l *is even,* $l \geq 2$,

$$[\mathcal{P}^{(s+4)}_{0,1,0,k-2}][\mathcal{P}^{(s)}_{0,1,0,k}] = [\mathcal{T}^{(s+8)}_{0,0,0,k-2}][\mathcal{T}^{(s+4)}_{0,0,0,k}][\mathcal{P}^{(s)}_{0,2,0,k-2}] + [\mathcal{P}^{(s)}_{0,0,2,k}][\mathcal{T}^{(s+4)}_{1,0,0,k-2}], \ k \geq 3, \tag{5.19}$$

$$[\mathcal{P}^{(s+4)}_{0,m,0,k-2}][\mathcal{P}^{(s)}_{0,m,0,k}] = [\mathcal{P}^{(s+4)}_{0,m-1,0,k}][\mathcal{P}^{(s)}_{0,m+1,0,k-2}] + [\mathcal{P}^{(s)}_{0,0,2m,k}][\mathcal{T}^{(s+4)}_{m,0,0,k-2}], \ k \geq 3, \ m \geq 2, \tag{5.20}$$

$$[\mathcal{T}^{(s+4)}_{n,0,0,k-2}][\mathcal{T}^{(s)}_{n,0,0,k}] = [\mathcal{T}^{(s+4)}_{n-1,0,0,k}][\mathcal{T}^{(s)}_{n+1,0,0,k-2}] + [\mathcal{P}^{(s)}_{0,n,0,k}], \ k \geq 3. \tag{5.21}$$

Moreover, every module in the summands on the right hand side of above equation is simple.

Proof. The lowest weight monomial of $\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$ is obtained from the highest weight monomial of $\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$ by the substitutions: $1_s \mapsto 1^{-1}_{18+s}$, $2_s \mapsto 2^{-1}_{18+s}$, $3_s \mapsto 3^{-1}_{18+s}$, $4_s \mapsto 4^{-1}_{18+s}$. After we apply ι to $\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$, the lowest weight monomial of $\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$ becomes the highest weight monomial of $\iota(\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}))$. Therefore the highest weight monomial of $\iota(\chi_q(\mathcal{T}^{(s)}_{n,m,l,k}))$ is obtained from the lowest weight monomial of $\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$ by the substitutions: $1_s \mapsto 1^{-1}_{18-s}$, $2_s \mapsto 2^{-1}_{18-s}$, $3_s \mapsto 3^{-1}_{18-s}$, $4_s \mapsto 4^{-1}_{18-s}$. It follows that the highest weight monomial of $\iota(\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}))$ is obtained from the highest weight monomial of $\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$ by the substitutions: $1_s \mapsto 1_{-s}$, $2_s \mapsto 2_{-s}$, $3_s \mapsto 3_{-s}$, $4_s \mapsto 4_{-s}$. Therefore the dual system is obtained applying ι to both sides of every equation of the system in Theorem 3.4.

The simplify of every module in the summands on the right hand side of every equation follows from Theorem 3.6 and Lemma 5.3. $\square$

Example 5.5. *The following are some equations in the system in Theorem 5.4.*

$$
\begin{aligned}
&[1_2][1_4 2_1] = [1_4 1_2][2_1] + [2_3 2_1],\\
&[1_4 1_2][1_6 1_4 2_1] = [1_4 2_1][1_6 1_4 1_2] + [2_5 2_3 2_1],\\
&[1_6 1_4 1_2][1_8 1_6 1_4 2_1] = [1_6 1_4 2_1][1_8 1_6 1_4 1_2] + [2_7 2_5 2_3 2_1],\\
&[3_4][1_0 2_3 3_8] = [1_0 2_3][3_4 3_8] + [1_0 2_3 2_5 2_7][4_6],\\
&[3_4 3_8][1_0 2_3 3_8 3_{12} = [1_0 2_3 3_8][3_4 3_8 3_{12}] + [1_0 2_3 2_5 2_7 2_9 2_{11}][4_6 4_{10}],\\
&[3_4 3_8 3_{12}][1_0 2_3 3_8 3_{12} 3_{16}] = [1_0 2_3 3_8 3_{12}][3_4 3_8 3_{12} 3_{16}] + [1_0 2_3 2_5 2_7 2_9 2_{11} 2_{13} 2_{15}][4_6 4_{10} 4_{14}],\\
&[4_6][1_0 2_3 4_{10}] = [1_0 2_3][4_6 4_{10}] + [1_0 2_3 3_8],\\
&[4_6 4_{10}][1_0 2_3 4_{10} 4_{14}] = [1_0 2_3 4_{10}][4_6 4_{10} 4_{14}] + [1_0 2_3 3_8 3_{12}],\\
&[4_6 4_{10} 4_{14}][1_0 2_3 4_{10} 4_{14} 4_{18}] = [1_0 2_3 4_{10} 4_{14}][4_6 4_{10} 4_{14} 4_{18}] + [1_0 2_3 3_8 3_{12} 3_{16}].
\end{aligned}
$$

5.1. **The system in Theorem 5.4.** By replacing each $[\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}]$ (resp. $[\mathcal{T}^{(s)}_{n,m,l,k}]$) in the system of Theorem 5.4 with $\chi(\widetilde{\mathfrak{m}}_{n,m,l,k})$ (resp. $\chi(\mathfrak{m}_{n,m,l,k})$), we obtain a system of equations consisting of the characters of $U_q\mathfrak{g}$-modules. The following are two equations in the system.

$$
\begin{aligned}
&\chi(\widetilde{\mathfrak{m}}_{0,0,l-1,k})\chi(\widetilde{\mathfrak{m}}_{0,0,l,k}) = \chi(\widetilde{\mathfrak{m}}_{0,0,l,k-1})\chi(\widetilde{\mathfrak{m}}_{0,0,l-1,k+1}) + \chi(\widetilde{\mathfrak{m}}_{0,0,k+l,0})\chi(\widetilde{\mathfrak{m}}_{0,0,l-1,0}),\\
&\chi(\mathfrak{m}_{n,m-1,0,0})\chi(\mathfrak{m}_{n,m,0,0}) = \chi(\mathfrak{m}_{n-1,m,0,0})\chi(\mathfrak{m}_{n+1,m-1,0,0}) + \chi(\mathfrak{m}_{0,m-1,0,0})\chi(\mathfrak{m}_{0,n+m,0,0}).
\end{aligned}
$$

5.2. **Relation between the system in Theorem 5.4 and cluster algebras.** Let $I = \{1, 2, 3, 4\}$ and

$$
\begin{aligned}
S &= \{2u \mid u \in \mathbb{Z}_{\geq 0}\},\\
S' &= \{2u + 1 \mid u \in \mathbb{Z}_{\geq 0}\}.
\end{aligned}
$$

Let

$$V = (\{1\} \times S) \cup (\{2\} \times S') \cup (\{3\} \times S) \cup (\{4\} \times S).$$

A quiver $\widetilde{Q}$ with vertex set V will be defined as follows. The arrows of $\widetilde{Q}$ are given by the following rule: there is an arrow from the vertex (i, r) to the vertex (j, s) if and only if $b_{ij} \neq 0$ and $s = r - b_{ij} + d_i - d_j$.

Let $\widetilde{\mathbf{t}} = \widetilde{\mathbf{t}}_1 \cup \widetilde{\mathbf{t}}_2$, where

$$\widetilde{\mathbf{t}}_{\mathbf{1}} = \{\widetilde{t}^{(-2)}_{0,0,l,0},\ \widetilde{t}^{(-4)}_{0,0,0,k} \mid k, l \in \mathbb{Z}_{\geq 1}\}$$

and

$$\widetilde{\mathbf{t}}_2 = \{t^{(-4n+4)}_{n,0,0,0},\ t^{(-4m)}_{0,m,0,0},\ t^{(-4n+2)}_{n,0,0,0},\ t^{(-4m+2)}_{0,m,0,0},\ t^{(-2l-2)}_{0,0,l,0},\ t^{(2k-2)}_{0,0,0,k} \mid k, l, m, n \in \mathbb{Z}_{\geq 1}\}.$$

Let $\widetilde{\mathscr{A}}$ be the cluster algebra defined by the initial seed $(\widetilde{\mathbf{t}}, \widetilde{Q})$. By similar arguments in Section 4, we have the following theorem.

Theorem 5.6. *Every equation in the system in Theorem 5.4 corresponds to a mutation equation in the cluster algebra $\widetilde{\mathscr{A}}$. Every minimal affinization in the system in Theorem 5.4 corresponds to a cluster variable of the cluster algebra $\widetilde{\mathscr{A}}$.*

6. Proof of theorem 3.3

In this section, we prove Theorem 3.3. Namely, we will prove that for $s \in \mathbb{Z}, k, l, m, n \in \mathbb{Z}_{\geq 1}$, the modules

$$\begin{gathered}\mathcal{T}^{(-2)}_{0,0,l,k},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{0,m,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,0,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,0,k},\ \widetilde{\mathcal{T}}^{(s)}_{n,m,0,k}\ (k \leq 2),\\ \widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}\ (k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{n,0,l,k}\ (k \leq 2),\ \widetilde{\mathcal{T}}^{(s)}_{0,m,l,k}\ (k \leq 2),\ \widetilde{P}^{(s)}_{0,0,l,k},\ \widetilde{P}^{(s)}_{0,m,0,k},\end{gathered} \tag{6.1}$$

are special. Since the modules

$$\mathcal{T}^{(s)}_{0,0,0,k},\ \mathcal{T}^{(s)}_{0,0,l,0},\ \mathcal{T}^{(s)}_{0,m,0,0},\ \mathcal{T}^{(s)}_{m,0,0,0},\ \widetilde{\mathcal{T}}^{(s)}_{0,0,0,k},\ \widetilde{\mathcal{T}}^{(s)}_{0,0,l,0},\ \widetilde{\mathcal{T}}^{(s)}_{0,m,0,0},\ \widetilde{\mathcal{T}}^{(s)}_{n,0,0,0},$$

are Kirillov-Reshetikhin modules, they are special. By Theorem 3.1, the modules $\widetilde{\mathcal{T}}^{(s)}_{n,m,0,0}$, $\widetilde{\mathcal{T}}^{(s)}_{n,0,l,0}$, $\widetilde{\mathcal{T}}^{(s)}_{0,m,l,0}$, $\widetilde{\mathcal{T}}^{(s)}_{n,m,l,0}$ are special. In the following, we will prove that the other modules in (6.1) are special. Without loss of generality, we may assume that $s = 0$ in $\widetilde{\mathcal{T}}^{(s)}$, where $\widetilde{\mathcal{T}}$ is a module in (6.1).

6.1. **The cases of $\widetilde{\mathcal{T}}^{(0)}_{n,0,0,k}$, $\mathcal{T}^{(-2)}_{0,0,l,k}$, $\widetilde{P}^{(s)}_{0,0,l,k}$, and $\widetilde{P}^{(s)}_{0,m,0,k}$.** Since the proof of each case is similar to each other, we give a detailed proof of the case of $\widetilde{\mathcal{T}}^{(0)}_{n,0,0,k}$ as follows.

Let $m_+ = \widetilde{T}^{(0)}_{n,0,0,k}$ with $n, k \in \mathbb{Z}_{\geq 1}$. Then

$$m_+ = (4_0 4_4 \cdots 4_{4n-4})(1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2}).$$

Suppose that $n = 1$. Let

$$U = I \times \{aq^s : s \in \mathbb{Z}, s \leq 2k + 6\}.$$

Since all monomials in $\mathscr{M}(\chi_q(m_+) - \operatorname{trunc}_{m_+\mathcal{Q}_U^-} \chi_q(m_+))$ are right-negative, it is sufficient to show that $\operatorname{trunc}_{m_+\mathcal{Q}_U^-} \chi_q(m_+)$ is special.

Let

$$\mathscr{M} = \{m_0 = m_+, m_1 = m_0 A^{-1}_{4,2}, m_2 = m_1 A^{-1}_{3,4}, m_3 = m_2 A^{-1}_{2,6}, m_4 = m_3 A^{-1}_{2,4}, m_5 = m_4 A^{-1}_{3,6}, m_6 = m_5 A^{-1}_{4,8}\}.$$

It is easy to see that $\mathscr{M}$ satisfies the conditions in Theorem 2.4. Therefore

$$\operatorname{trunc}_{m_+\mathcal{Q}_U^-} \chi_q(m_+) = \sum_{m \in \mathscr{M}} m$$

and hence $\operatorname{trunc}_{m_+\mathcal{Q}_U^-}\chi_q(m_+)$ is special.

Suppose that $n \geq 2$. In the following, we write $m_+ = m_1'm_2' = m_1''m_2''$ for some monomials m_1', m_2', m_1'', m_2''. We will show that the only dominant monomial in $\mathscr{M}(\chi_q(m_1')\chi_q(m_2')) \cap \mathscr{M}(\chi_q(m_1'')\chi_q(m_2''))$ is m_+ which implies that $L(m_+)$ is special.

Let $m_+ = m_1'm_2'$, where

$$m_1' = 4_0 4_4 \cdots 4_{4n-8},\ m_2' = 4_{4n-4} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2}.$$

We have shown that $L(m_2')$ is special. Therefore the Frenkel-Mukhin algorithm works for $L(m_2')$. We will use the Frenkel-Mukhin algorithm to compute $\chi_q(L(m_1')), \chi_q(L(m_2'))$ and classify all dominant monomials in $\chi_q(L(m_1'))\chi_q(L(m_2'))$. Let $m = m_1m_2$ be a dominant monomial, where $m_i \in \mathscr{M}(L(m_i'))$, $i = 1, 2$.

Suppose that $m_2 \neq m_2'$. If m_2 is right-negative, then m is a right negative monomial and therefore m is not dominant, this is a contradiction. Hence m_2 is not right-negative. Through the above discussion, m_2 is one of the following monomials

$$\begin{aligned}
\bar{m}_1 &= m_2'A_{4,4n-2}^{-1} = 4_{4n}^{-1} 3_{4n-2} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2},\\
\bar{m}_2 &= \bar{m}_1 A_{3,4n}^{-1} = 3_{4n+2}^{-1} 2_{4n-1} 2_{4n+1} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2},\\
\bar{m}_3 &= \bar{m}_2 A_{2,4n+2}^{-1} = 2_{4n-1} 2_{4n+3}^{-1} 1_{4n+2} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2},\\
\bar{m}_4 &= \bar{m}_2 A_{2,4n}^{-1} = 3_{4n} 2_{4n+1}^{-1} 2_{4n+3}^{-1} 1_{4n} 1_{4n+2} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2},\\
\bar{m}_5 &= \bar{m}_4 A_{3,4n+2}^{-1} = 4_{4n+2} 3_{4n+4}^{-1} 1_{4n} 1_{4n+2} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2},\\
\bar{m}_6 &= \bar{m}_4 A_{4,4n+4}^{-1} = 4_{4n+6}^{-1} 1_{4n} 1_{4n+2} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2}.
\end{aligned}$$

We nextly discuss m_2 as follows:

Case 1. The factor 4_{4n} can only come from the monomials in $\chi_q(4_{4n-8})$, the monomials in $\chi_q(4_{4n-8})$ which contain a factor 4_{4n} are

$$\begin{aligned}
&3_{4n+2}^{-1} 4_{4n-2} 4_{4n},\ 2_{4n+1} 2_{4n+3} 3_{4n+2}^{-1} 3_{4n+4}^{-1} 4_{4n},\ 1_{4n+4} 2_{4n+1} 2_{4n+5}^{-1} 3_{4n+2}^{-1} 4_{4n},\ 1_{4n+4}^{-1} 1_{4n+6}^{-1} 4_{4n},\\
&1_{4n+2} 1_{4n+4} 2_{4n+3}^{-1} 2_{4n+5}^{-1} 4_{4n},\ 1_{4n+6}^{-1} 2_{4n+1} 3_{4n+2}^{-1} 4_{4n},\ 1_{4n+2} 1_{4n+6}^{-1} 2_{4n+3}^{-1} 4_{4n}.
\end{aligned} \tag{6.2}$$

The negative factors in (6.2) can not be canceled by monomials in $\chi_q(4_{4n-12})$. However, the negative factor $1_{4n+4}^{-1} 1_{4n+6}^{-1} 4_{4n}$ can be canceled by $\bar{m}_1$. Therefore $m_2 = \bar{m}_1$ in the situation.

Case 2. The factor 3_{4n+2} can only come from the monomials in $\chi_q(4_{4n-8})$, the monomials in $\chi_q(4_{4n-8})$ which contain a factor 3_{4n+2} are

$$1_{4n+2} 1_{4n+4} 2_{4n+3}^{-1} 2_{4n+5}^{-1} 3_{4n+2} 4_{4n+4}^{-1},\ 1_{4n+2} 1_{4n+6}^{-1} 2_{4n+3}^{-1} 3_{4n+2} 4_{4n+4}^{-1},\ 1_{4n+4}^{-1} 1_{4n+6}^{-1} 3_{4n+2} 4_{4n+4}^{-1}. \tag{6.3}$$

The negative factors in (6.3) can not be canceled by monomials in $\chi_q(4_{4n-12})$.

Case 3. The factors $2_{4n+3}, 3_{4n+4}, 4_{4n+6}$ can only come from the monomials in $\chi_q(4_{4n-8})$, the monomials in $\chi_q(4_{4n-8})$ which contain factors $2_{4n+3}, 3_{4n+4}, 4_{4n+6}$ are

$$\begin{aligned}
&1_{4n-4} 2_{4n+3} 3_{4n+4}^{-1},\quad 1_{4n-2}^{-1} 2_{4n-3} 2_{4n+3} 3_{4n+4}^{-1},\quad 2_{4n-1}^{-1} 2_{4n+3} 3_{4n-2} 3_{4n+4}^{-1},\\
&2_{4n+1} 2_{4n+3} 3_{4n+2}^{-1} 3_{4n+4}^{-1} 4_{4n},\quad 2_{4n+1} 2_{4n+3} 3_{4n+4}^{-1} 4_{4n+4}^{-1},\quad 1_{4n+4}^{-1} 2_{4n+3} 2_{4n+7}^{-1},\\
&1_{4n+4}^{-1} 1_{4n+6}^{-1} 2_{4n+3} 2_{4n+5} 3_{4n+6}^{-1},\quad 3_{4n+8}^{-1} 4_{4n+6},\quad 2_{4n+5}^{-1} 2_{4n+7}^{-1} 3_{4n+4}.
\end{aligned} \tag{6.4}$$

The negative factors in (6.4) can not be canceled by monomials in $\chi_q(4_{4n-12})$.

Since

$$\chi_q(4_0 4_4 \cdots 4_{4n-8}) \subseteq \chi_q(4_0 4_4 \cdots 4_{4n-12})\chi_q(4_{4n-8}) \subseteq \chi_q(4_0 4_4 \cdots 4_{4n-16})\chi_q(4_{4n-12})\chi_q(4_{4n-8}),$$

and $m = m_1 m_2$ is dominant, $m_2 = m'_2$ or $\bar{m}_1$.

Suppose that $m_2 = \bar{m}_1$. By the above discussion, m is in

$$\chi_q(4_0 4_4 \cdots 4_{4n-12}) 3_{4n-2} 1_{4n+8} \cdots 1_{4n+2k+2}. \tag{6.5}$$

Suppose that $m_2 = m'_2$. If $m_1 \neq m'_1$, then m_1 is right negative. Since m is dominant, each factor with a negative power in m_1 needs to be canceled by a factor in m'_2. We have $\mathscr{M}(L(m'_1)) \subset \mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12}))\ \chi_q(L(4_{4n-8}))$. Only monomials in $\chi_q(L(4_{4n-8}))$ can cancel 4_{4n-4}, 1_{4n+4}, 1_{4n+6}. Therefore m_1 is one of the sets

$$\begin{aligned}
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 3_{4n-6} 4^{-1}_{4n-4},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+4} 1^{-1}_{4n+6} 4_{4n},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+4} 1^{-1}_{4n+6} 3_{4n+2} 4_{4n+4},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+4} 1^{-1}_{4n+6} 2_{4n+3} 2_{4n+5} 3^{-1}_{4n+6},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+4} 2_{4n+3} 2^{-1}_{4n+7},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1_{4n-4} 1^{-1}_{4n+6},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n-2} 1^{-1}_{4n+6} 2_{4n-3},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+6} 2^{-1}_{4n-1} 3_{4n-2},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+6} 2_{4n+1} 3^{-1}_{4n+2} 4_{4n},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1_{4n+2} 1^{-1}_{4n+6} 2^{-1}_{4n+3} 4_{4n},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+6} 2_{4n+1} 4^{-1}_{4n+4},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1_{4n+2} 1^{-1}_{4n+6} 2^{-1}_{2n+3} 3_{4n+2} 4^{-1}_{4n+4},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1_{4n+2} 1^{-1}_{4n+6} 2_{4n+5} 3^{-1}_{4n+6}.
\end{aligned}$$

By (6.4), we know that $m_1 \notin \mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1^{-1}_{4n+6} 2^{-1}_{4n-1} 3_{4n-2}$. The factors which contain 1_{4n-2} in $\chi_q(4n-12)$ are

$$\begin{gathered}
1_{4n-2} 1_{4n} 2^{-1}_{4n-1} 2^{-1}_{4n+1} 4_{4n-4},\ 1_{4n-2} 1_{4n} 2^{-1}_{4n-1} 2^{-1}_{4n+1} 3_{4n-2} 4^{-1}_{4n},\ 1_{4n-2} 1^{-1}_{4n+2} 2^{-1}_{4n-1} 4_{4n-4},\\
1_{4n-2} 1^{-1}_{4n+2} 2^{-1}_{4n-1} 3_{4n-2} 4^{-1}_{4n},\ 1_{4n-2} 1_{4n} 3^{-1}_{4n+2},\ 1_{4n-2} 1^{-1}_{4n+2} 2_{4n+1} 3^{-1}_{4n+2},\ 1_{4n-2} 2^{-1}_{4n+3}.
\end{gathered} \tag{6.6}$$

The negative factors in (6.6) can not be canceled by monomials in $\chi_q(4_{4n-16})$.

Since $\chi_q(4_0 4_4 \cdots 4_{4n-12}) \subseteq \chi_q(4_0 4_4 \cdots 4_{4n-16})\chi_q(4n-12)$ and 1_{4n+4}, 1_{4n+6}, $2_{4n+7} \notin \chi_q(4n-12)$. Therefore m_1 is in one of the sets

$$\begin{aligned}
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 3_{4n-6} 4^{-1}_{4n-4},\\
&\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1_{4n-4} 1^{-1}_{4n+6}.
\end{aligned}$$

Therefore m is in one of the sets

$$\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 3_{4n-6} 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2}, \tag{6.7}$$

$$\mathscr{M}(\chi_q(4_0 4_4 \cdots 4_{4n-12})) 1_{4n-4} 4_{4n-4} 1_{4n+4} 1_{4n+8} \cdots 1_{4n+2k+2}. \tag{6.8}$$

Let $m_+ = m_1'' m_2''$, where

$$m_1'' = 4_0 4_4 \cdots 4_{4n-4}, \;\; m_2'' = 1_{4n+4} 1_{4n+6} \cdots 1_{4n+2k+2}.$$

If m is the expressions of (6.5) (6.7), (6.8), we know that $m \notin \mathscr{M}(\chi_q(m_1'')\chi_q(m_2''))$ by the Frenkel-Mukhin algorithm.

Therefore the only dominant monomial in $\mathscr{M}(\chi_q(m_1')\chi_q(m_2')) \cap \mathscr{M}(\chi_q(m_1'')\chi_q(m_2''))$ is m_+. Hence the only dominant monomial in $\chi_q(m_+)$ is m_+.

6.2. The case of $\widetilde{\mathcal{T}}^{(0)}_{n,m,0,k}$ $(k \leq 2)$, $\widetilde{\mathcal{T}}^{(0)}_{n,0,l,k}$ $(k \leq 2)$, and $\widetilde{\mathcal{T}}^{(0)}_{0,m,l,k}$ $(k \leq 2)$.

Since the proof of each case is similar to each other, we give a detailed proof of the case of $\widetilde{\mathcal{T}}^{(0)}_{n,m,0,k}$ $(k \leq 2)$ as follows.

Let $m_+ = \widetilde{T}^{(0)}_{n,m,0,k}$ with $n, m, k \in \mathbb{Z}_{\geq 1}$ and $k \leq 2$. Then

$$m_+ = (4_0 4_4 \cdots 4_{4n-4})(3_{4n+2} 3_{4n+6} \cdots 3_{4n+4m-2})(1_{4n+4m+4} \cdots 1_{4n+4m+2k+2}).$$

Let

$$\begin{aligned} m_1' &= (4_0 4_4 \cdots 4_{4n-4})(3_{4n+2} 3_{4n+6} \cdots 3_{4n+4m-2}), \;\; m_2' = (1_{4n+4m+4} \cdots 1_{4n+4m+2k+2}), \\ m_1'' &= (4_0 4_4 \cdots 4_{4n-4}), \;\; m_2'' = (3_{4n+2} 3_{4n+6} \cdots 3_{4n+4m-2})(1_{4n+4m+4} \cdots 1_{4n+4m+2k+2}). \end{aligned}$$

Then $\mathscr{M}(L(m_+)) \subset \mathscr{M}(\chi_q(m_1')\chi_q(m_2')) \cap \mathscr{M}(\chi_q(m_1'')\chi_q(m_2''))$.

By using similar arguments as in Subsection 6.1, we can show that the only possible dominant monomial in $\chi_q(m_1')\chi_q(m_2') \cap \chi_q(m_1'')\chi_q(m_2'')$ is m_+. Hence the only dominant monomial in $\chi_q(m_+)$ is m_+.

6.3. The case of $\widetilde{\mathcal{T}}^{(0)}_{n,m,l,k}(k \leq 2)$.

Let $m_+ = \widetilde{\mathcal{T}}^{(0)}_{n,m,l,k}$ with $n, m, l, k \in \mathbb{Z}_{\geq 1}$ and $k \leq 2$. Then

$$\begin{aligned} m_+ =& (4_0 4_4 \cdots 4_{4n-4})(3_{4n+2} 3_{4n+6} \cdots 3_{4n+4m-2})(2_{4n+4m+3} 2_{4n+4m+5} \cdots 2_{4n+4m+2l+1}) \\ & (1_{4n+4m+2l+4} 1_{4n+4m+2l+1+6} \cdots 1_{4n+4m+2l+2k+2}). \end{aligned}$$

Let

$$\begin{aligned} m_1' &= (4_0 4_4 \cdots 4_{4n-4})(3_{4n+2} 3_{4n+6} \cdots 3_{4n+4m-2})(2_{4n+4m+3} 2_{4n+4m+5} \cdots 2_{4n+4m+2l+1}), \\ m_2' &= (1_{4n+4m+2l+4} 1_{4n+4m+2l+1+6} \cdots 1_{4n+4m+2l+2k+2}), \\ m_1'' &= (4_0 4_4 \cdots 4_{4n-4}), \\ m_2'' &= (3_{4n+2} 3_{4n+6} \cdots 3_{4n+4m-2})(2_{4n+4m+3} 2_{4n+4m+5} \cdots 2_{4n+4m+2l+1})(1_{4n+4m+2l+4} 1_{4n+4m+2l+1+6} \cdots 1_{4n+4m+2l+2k+2}). \end{aligned}$$

Then $\mathscr{M}(L(m_+)) \subset \mathscr{M}(\chi_q(m_1')\chi_q(m_2')) \cap \mathscr{M}(\chi_q(m_1'')\chi_q(m_2''))$.

By using similar arguments as in Subsection 6.1, we can show that the only possible dominant monomial in $\chi_q(m_1')\chi_q(m_2') \cap \chi_q(m_1'')\chi_q(m_2'')$ is m_+. Hence the only dominant monomial in $\chi_q(m_+)$ is m_+.

7. PROOF OF THEOREM 3.4

In this section, we prove Theorem 3.4.

7.1. **Classification of dominant monomials in the summands on both sides of the system.** In Section 6, we have shown that for $s \in \mathbb{Z}, k, l, m, n \in \mathbb{Z}_{\geq 1}$, the modules in Theorem 3.3 are special. Now we use the Frenkel-Mukhin algorithm to classify dominant monomials in the summands on both sides of the system in Theorem 3.4.

Lemma 7.1. *The dominant monomials in each summand on the left and right hand sides of every equation in the system of Theorem 3.4 are given in Table 2 and 3.*

Proof. We will prove the case of (3.15), the other cases are similar.

Let $m_1' = \widetilde{T}^{(s+4)}_{n,m-1,l,k}$, $m_2' = \widetilde{T}^{(s)}_{n,m,l,k}$. Without loss of generality, we may assume that $s = 0$. Then

$$\begin{aligned} m_1' =& (4_4 \cdots 4_{4n})(3_{4n+6}3_{4n+10} \cdots 3_{4n+4m-2})(2_{4n+4m+3}2_{4n+4m+5} \cdots 2_{4n+4m+2l+1}) \\ & (1_{4n+4m+2l+4}1_{4n+4m+2l+6} \cdots 1_{4n+4m+2l+2k+2}), \\ m_2' =& (4_0 4_4 \cdots 4_{4n-4})(3_{4n+2}3_{4n+6} \cdots 3_{4n+4m-2})(2_{4n+4m+3}2_{4n+4m+5} \cdots 2_{4n+4m+2l+1}) \\ & (1_{4n+4m+2l+4}1_{4n+4m+2l+6} \cdots 1_{4n+4m+2l+2k+2}). \end{aligned}$$

Let $m = m_1 m_2$ be a dominant monomial, where $m_i \in \chi_q(m_i'), i = 1, 2$. We denote

$$\begin{aligned} m_3 =& (3_{4n+6}3_{4n+10} \cdots 3_{4n+4m-2})(2_{4n+4m+3}2_{4n+4m+5} \cdots 2_{4n+4m+2l+1}) \\ & (1_{4n+4m+2l+4}1_{4n+4m+2l+6} \cdots 1_{4n+4m+2l+2k+2}), \\ m_4 =& (3_{4n+2}3_{4n+6} \cdots 3_{4n+4m-2})(2_{4n+4m+3}2_{4n+4m+5} \cdots 2_{4n+4m+2l+1}) \\ & (1_{4n+4m+2l+4}1_{4n+4m+2l+6} \cdots 1_{4n+4m+2l+2k+2}). \end{aligned}$$

Suppose that $m_1 \in \chi_q(4_4 \cdots 4_{4n})(\chi_q(m_3) - m_3)$, then $m = m_1 m_2$ is right negative and hence m is not dominant. Therefor $m_1 \in \chi_q(4_4 \cdots 4_{4n})m_3$. Similarly, if $m_2 \in \chi_q(4_0 4_4 \cdots 4_{4n-4})(\chi_q(m_4) - m_4)$, then $m = m_1 m_2$ is right negative and hence m is not dominant. This contradicts our assumption. Therefore $m_2 \in \chi_q(4_0 4_4 \cdots 4_{4n-4})m_4$.

Suppose that $m_1 \in \mathscr{M}(L(m_1')) \cap \mathscr{M}(\chi_q(4_4 \cdots 4_{4n-4})(\chi_q(4_{4n}) - 4_{4n})m_3)$. By the Frenkel-Mukhin algorithm for $L(m_1')$, m_1 must have the factor 4^{-1}_{4n+4}. But by the Frenkel-Mukhin algorithm and the fact that $m_2 \in \chi_q(4_0 4_4 \cdots 4_{4n-4})m_4$, m_2 does not have the factor 4_{4n+4}. Therefore $m_1 m_2$ is not dominant. Hence $m_1 \in \chi_q(4_4 \cdots 4_{4n-4})4_{4n}m_3$. It follows that $m_1 = m_1'$.

By the Frenkel-Mukhin algorithm and the fact that $m_2 \in \chi_q(4_0 4_4 \cdots 4_{4n-4})m_4$, m_2 must be one of the following monomials,

$$\begin{aligned} v_1 &= m_2' A^{-1}_{4,4n-2} = 4_0 4_4 \cdots 4_{4n-8} 4^{-1}_{4n} 3_{4n-2} m_4, \\ v_2 &= m_2' A^{-1}_{4,4n-2} A^{-1}_{4,4n-6} = 4_0 4_4 \cdots 4_{4n-12} 4^{-1}_{4n-4} 4^{-1}_{4n} 3_{4n-6} 3_{4n-2} m_4, \\ & \cdots \\ v_n &= m_2' A^{-1}_{4,4n-2} A^{-1}_{4,4n-6} \cdots A^{-1}_{4,2} = 4^{-1}_4 \cdots 4^{-1}_{4n-4} 4^{-1}_{4n} 3_2 \cdots 3_{4n-6} 3_{4n-2} m_4. \end{aligned}$$

It follows that the dominant monomials in $\chi_q(\widetilde{T}^{(s+4)}_{n,m-1,l,k})\chi_q(\widetilde{T}^{(s)}_{n,m,l,k})$ are

Equations	Summands in the equations	M	Dominant monomials
(3.1)	$\chi_q(\mathcal{T}^{(-2)}_{0,0,l-1,k})\chi_q(\mathcal{T}^{(-2)}_{0,0,l,k})$	$M=T^{(-2)}_{0,0,l-1,k}T^{(-2)}_{0,0,l,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{1,aq^{-2l-1-2i}}$, $0\le r\le k$
(3.1)	$\chi_q(\mathcal{T}^{(-2)}_{0,0,l,k-1})\chi_q(\mathcal{T}^{(-2)}_{0,0,l-1,k+1})$	$M=T^{(-2)}_{0,0,l,k-1}T^{(-2)}_{0,0,l-1,k+1}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{1,aq^{-2l-1-2i}}$, $0\le r\le k-1$
(3.1)	$\chi_q(\mathcal{T}^{(-2)}_{0,0,k+l,0})\chi_q(\mathcal{T}^{(-2)}_{0,0,l-1,0})$	$M=T^{(-2)}_{0,0,k+l,0}T^{(-2)}_{0,0,l-1,0}$	M
(3.2)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,0,0})$	$M=\widetilde{T}^{(s+4)}_{n,m-1,0,0}\widetilde{T}^{(s)}_{n,m,0,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.2)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,m+1,0,0})$	$M=\widetilde{T}^{(s+4)}_{n-1,m,0,0}\widetilde{T}^{(s)}_{n+1,m-1,0,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $-1\le r\le n-1$
(3.2)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,m-1,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n+m,0,0})$	$M=\widetilde{T}^{(s+4n+4)}_{0,0,l-2,0}\widetilde{T}^{(s)}_{0,n,l,0}$	M
(3.3)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,1,0})$	$M=\widetilde{T}^{(s+4)}_{n,0,0,0}\widetilde{T}^{(s)}_{n,0,1,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.3)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,1,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,0})$	$M=\widetilde{T}^{(s+4)}_{n-1,0,1,0}\widetilde{T}^{(s)}_{n+1,0,0,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $-1\le r\le n-1$
(3.3)	$\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n,l,0})$	$M=\widetilde{T}^{(s)}_{0,n,l,0}$	M
(3.4)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,l-2,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,l,0})$	$M=\widetilde{T}^{(s+4)}_{n,0,l-2,0}\widetilde{T}^{(s)}_{n,0,l,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.4)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,l,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,0,l-2,0})$	$M=\widetilde{T}^{(s+4)}_{n-1,0,l,0}\widetilde{T}^{(s)}_{n+1,0,l-2,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $-1\le r\le n-1$
(3.4)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,0,l-2,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n,l,0})$	$M=\widetilde{T}^{(s+4n+4)}_{0,0,l-2,0}\widetilde{T}^{(s)}_{0,n,l,0}$	M
(3.5)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,1,0})$	$M=\widetilde{T}^{(s+4)}_{0,m,0,0}\widetilde{T}^{(s)}_{0,m,1,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m$
(3.5)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,1,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,0})$	$M=\widetilde{T}^{(s+4)}_{0,m,0,0}\widetilde{T}^{(s)}_{0,m,1,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m-1$
(3.5)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,0,1+2m,0})$	$M=\widetilde{T}^{(s+4)}_{m,0,0,0}\widetilde{T}^{(s)}_{0,0,1+2m,0}$	M
(3.6)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,l-2,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,l,0})$	$M=\widetilde{T}^{(s+4)}_{0,m,l-2,0}\widetilde{T}^{(s)}_{0,m,l,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m$
(3.6)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,l,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m+1,l-2,0})$	$M=\widetilde{T}^{(s+4)}_{0,m,l-2,0}\widetilde{T}^{(s)}_{0,m,l,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m-1$
(3.6)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,l-2})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,0,l+2m,0})$	$M=\widetilde{T}^{(s+4)}_{m,0,0,l-2}\widetilde{T}^{(s)}_{0,0,l+2m,0}$	M
(3.7)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,m,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,0})$	$M=\widetilde{T}^{(s+4)}_{n,m-1,l,0}\widetilde{T}^{(s)}_{n,m,l,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.7)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,0})$	$M=\widetilde{T}^{(s+4)}_{n-1,m,l,0}\widetilde{T}^{(s)}_{n+1,m-1,l,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n-1$
(3.7)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,m-1,l,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n+m,l,0})$	$M=\widetilde{T}^{(s+4n+4)}_{0,m-1,l,0})\widetilde{T}^{(s)}_{0,n+m,l,0}$	M
(3.8)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,0,k})$	$M=\widetilde{T}^{(s+4)}_{n,0,0,0}\widetilde{T}^{(s)}_{n,0,0,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.8)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,0})$	$M=\widetilde{T}^{(s+4)}_{n-1,0,0,k}\widetilde{T}^{(s)}_{n+1,0,0,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n-1$
(3.8)	$\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n,0,k})$	$M=\widetilde{T}^{(s)}_{0,n,0,k}$	M
(3.9)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,0,k})$	$M=\widetilde{T}^{(s+4)}_{0,m,0,0}\widetilde{T}^{(s)}_{0,m,0,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m$
(3.9)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,0})$	$M=\widetilde{T}^{(s+4)}_{0,m-1,0,k}\widetilde{T}^{(s)}_{0,m+1,0,0}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m-1$
(3.9)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,0,2m,k})$	$M=\widetilde{T}^{(s+4)}_{m,0,0,0}\widetilde{T}^{(s)}_{0,0,2m,k}$	M
(3.10)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-1})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,1,k})$	$M=\widetilde{T}^{(s+4)}_{n,0,0,k-1}\widetilde{T}^{(s)}_{n,0,1,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.10)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,1,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,k-1})$	$M=\widetilde{T}^{(s+4)}_{n-1,0,1,k}\widetilde{T}^{(s)}_{n+1,0,0,k-1}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n-1$
(3.10)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,0,0,k-1})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n,1,k})$	$M=\widetilde{T}^{(s+4n+4)}_{0,0,0,k-1}\widetilde{T}^{(s)}_{0,n,l,k}$	M
(3.11)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,l-2,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,l,k})$	$M=\widetilde{T}^{(s+4)}_{n,0,l-2,k}\widetilde{T}^{(s)}_{n,0,l,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.11)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,0,l-2,k})$	$M=\widetilde{T}^{(s+4)}_{n-1,0,l,k}\widetilde{T}^{(s)}_{n+1,0,l-2,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n-1$
(3.11)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,0,l-2,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n,l,k})$	$M=\widetilde{T}^{(s+4n+4)}_{0,0,l-2,k}\widetilde{T}^{(s)}_{0,n,l,k}$	M
(3.12)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,k-1})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,1,k})$	$M=\widetilde{T}^{(s+4)}_{0,m,0,k-1}\widetilde{T}^{(s)}_{0,m,1,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m$
(3.12)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,1,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,k-1})$	$M=\widetilde{T}^{(s+4)}_{0,m-1,1,k}\widetilde{T}^{(s)}_{0,m+1,0,k-1}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m-1$
(3.12)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,k-1})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,0,1+2m,k})$	$M=\widetilde{T}^{(s+4)}_{m,0,0,k-1}\widetilde{T}^{(s)}_{0,0,1+2m,k}$	M
(3.13)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,l-2,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,l,k})$	$M=\widetilde{T}^{(s+4)}_{0,m,l-2,k}\widetilde{T}^{(s)}_{0,m,l,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m$
(3.13)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m+1,l-2,k})$	$M=\widetilde{T}^{(s+4)}_{0,m-1,l,k}\widetilde{T}^{(s)}_{0,m+1,l-2,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0\le r\le m-1$
(3.13)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{m,0,l-2,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,0,l+2m,k})$	$M=\widetilde{T}^{(s+4)}_{m,0,l-2,k}\widetilde{T}^{(s)}_{0,0,l+2m,k}$	M
(3.14)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,0,k})$	$M=\widetilde{T}^{(s+4)}_{n,m-1,0,k}\widetilde{T}^{(s)}_{n,m,0,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.14)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,0,k})$	$M=\widetilde{T}^{(s+4)}_{n-1,m,0,k}\widetilde{T}^{(s)}_{n+1,m-1,0,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n-1$
(3.14)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n+m,0,k})$	$M=\widetilde{T}^{(s+4)}_{0,m-1,0,k}\widetilde{T}^{(s)}_{0,n+m,0,k}$	M
(3.15)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$	$M=\widetilde{T}^{(s+4)}_{n,m-1,l,k}\widetilde{T}^{(s)}_{n,m,l,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n$
(3.15)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,k})$	$M=\widetilde{T}^{(s+4)}_{n-1,m,l,k}\widetilde{T}^{(s)}_{n+1,m-1,l,k}$	$M_r=M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0\le r\le n-1$
(3.15)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4n+4)}_{0,m-1,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,n+m,l,k})$	$M=\widetilde{T}^{(s)}_{0,n+m,l,k}\widetilde{T}^{(s+4n+4)}_{0,m-1,l,k}$	M

TABLE 2. Classification of dominant monomials in the system in Theorem 3.4.

Equations	Summands in the equations	M	Dominant monomials
(3.16)	$\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,0,1,k-1})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,1,k})$	$M{=}\widetilde{P}^{(s+2)}_{0,0,1,k-1}\widetilde{P}^{(s)}_{0,0,1,k}$	$M_0{=}M,M_1{=}MA^{-1}_{2,aq^{s+4}}$
(3.16)	$\chi_q(\mathcal{T}^{(s+10)}_{0,0,0,k-4})\chi_q(\mathcal{T}^{(s+6)}_{0,0,0,k-2})\chi_q(\mathcal{T}^{(s+2)}_{0,0,0,k})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,2,k-1})$	$M{=}T^{(s+10)}_{0,0,0,k-4}T^{(s+6)}_{0,0,0,k-2}T^{(s+2)}_{0,0,0,k}\widetilde{P}^{(s)}_{0,0,2,k-1}$	M
(3.16)	$\chi_q(\mathcal{T}^{(s+12)}_{0,0,0,k-5})\chi_q(\mathcal{T}^{(s+8)}_{0,0,0,k-3})\chi_q(\mathcal{T}^{(s+4)}_{0,0,0,k-1})\chi_q(\mathcal{T}^{(s)}_{0,0,0,k+1})\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,1,0,k-2})$	$M{=}T^{(s+12)}_{0,0,0,k-5}T^{(s+8)}_{0,0,0,k-3}T^{(s+4)}_{0,0,0,k-1}T^{(s)}_{0,0,0,k+1}\widetilde{P}^{(s+2)}_{0,1,0,k-2}$	M
(3.17)	$\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l,k-1})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,l,k})$	$M{=}\widetilde{P}^{(s+2)}_{0,0,l,k-1}\widetilde{P}^{(s)}_{0,0,l,k}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{2,aq^{s+2l+2-2i}}$, $0{\leq}r{\leq}l$
(3.17)	$\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l-1,k})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,l+1,k-1})$	$M{=}\widetilde{P}^{(s+2)}_{0,0,l-1,k}\widetilde{P}^{(s)}_{0,0,l+1,k-1}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{2,aq^{s+2l+2-2i}}$, $0{\leq}r{\leq}l{-}1$
(3.17)	$\chi_q(\mathcal{T}^{(s+2l+10)}_{0,0,0,k-5})\chi_q(\mathcal{T}^{(s)}_{0,0,0,k+l})\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,\frac{l+1}{2},0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s+4)}_{0,\frac{l-1}{2},0,k-1})$	$M{=}T^{(s+2l+10)}_{0,0,0,k-5}T^{(s)}_{0,0,0,k+l}\widetilde{P}^{(s+2)}_{0,\frac{l+1}{2},0,k-2}\widetilde{\mathcal{P}}^{(s+4)}_{0,\frac{l-1}{2},0,k-1}$	M
(3.18)	$\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l,k-1})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,l,k})$	$M{=}\widetilde{P}^{(s+2)}_{0,0,l,k-1}\widetilde{P}^{(s)}_{0,0,l,k}$	$M_r{=}M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{2,aq^{s+2l+2-2i}}$, $0{\leq}r{\leq}l$
(3.18)	$\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l-1,k})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,l+1,k-1})$	$M{=}\widetilde{P}^{(s+2)}_{0,0,l-1,k}\widetilde{P}^{(s)}_{0,0,l+1,k-1}$	$M_r{=}M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{2,aq^{s+2l+2-2i}}$, $0{\leq}r{\leq}l{-}1$
(3.18)	$\chi_q(\mathcal{T}^{(s+2l+10)}_{0,0,0,k-5})\chi_q(\mathcal{T}^{(s)}_{0,0,0,k+l})\chi_q(\widetilde{\mathcal{P}}^{(s+4)}_{0,\frac{l}{2},0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,\frac{l}{2},0,k-1})$	$M{=}T^{(s+2l+10)}_{0,0,0,k-5}T^{(s)}_{0,0,0,k+l}\widetilde{P}^{(s+4)}_{0,\frac{l}{2},0,k-2}\widetilde{\mathcal{P}}^{(s+2)}_{0,\frac{l}{2},0,k-1}$	M
3.19)	$\chi_q(\widetilde{\mathcal{P}}^{(s+4)}_{0,1,0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,1,0,k})$	$M{=}\widetilde{P}^{(s+4)}_{0,1,0,k-2}\widetilde{P}^{(s)}_{0,1,0,k}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0{\leq}r{\leq}m$
(3.19)	$\chi_q(\mathcal{T}^{(s+8)}_{0,0,0,k-2})\chi_q(\mathcal{T}^{(s+4)}_{0,0,0,k})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,2,0,k-2})$	$M{=}T^{(s+8)}_{0,0,0,k-2}T^{(s+4)}_{0,0,0,k}\widetilde{P}^{(s)}_{0,2,0,k-2}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0{\leq}r{\leq}m{-}1$
(3.19)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{1,0,0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,2,k})$	$M{=}\widetilde{T}^{(s+4)}_{1,0,0,k-2}\widetilde{P}^{(s)}_{0,0,2,k}$	M
(3.20)	$\chi_q(\widetilde{\mathcal{P}}^{(s+4)}_{0,m,0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,m,0,k})$	$M{=}\widetilde{P}^{(s+4)}_{0,1,0,k-2}\widetilde{P}^{(s)}_{0,1,0,k}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0{\leq}r{\leq}m$
(3.20)	$\chi_q(\widetilde{\mathcal{P}}^{(s+4)}_{0,m-1,0,k})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,m+1,0,k-2})$	$M{=}\widetilde{P}^{(s+4)}_{0,m-1,0,k}\widetilde{P}^{(s)}_{0,m+1,0,k-2}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{3,aq^{s+4m-4i}}$, $0{\leq}r{\leq}m{-}1$
(3.20)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{m,0,0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,2m,k})$	$M{=}\widetilde{T}^{(s+4)}_{m,0,0,k-2}\widetilde{P}^{(s)}_{0,0,2m,k}$	M
(3.21)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-2})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,0,k})$	$M{=}\widetilde{T}^{(s+4)}_{n,0,0,k-2}\widetilde{T}^{(s)}_{n,0,0,k}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0{\leq}r{\leq}n$
(3.21)	$\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,k-2})$	$M{=}\widetilde{T}^{(s+4)}_{n-1,0,0,k}\widetilde{T}^{(s)}_{n+1,0,0,k-2}$	$M_r{=}M\prod_{i=0}^{r-1}A^{-1}_{4,aq^{s+4n-2-4i}}$, $0{\leq}r{\leq}n{-}1$
(3.21)	$\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,n,0,k})$	$M{=}\widetilde{P}^{(s)}_{0,n,0,k}$	M

TABLE 3. Classification of dominant monomials in the system in Theorem 3.4 (continued).

$$M = m_1' m_2',\ M_1 = v_1 m_1' = M A^{-1}_{4,4n-2},\ M_2 = v_2 m_1' = M \prod_{i=0}^{1} A^{-1}_{4,4n-4i-2},\ \ldots,$$

$$M_{n-1} = v_{n-1} m_1' = M \prod_{i=0}^{n-2} A^{-1}_{4,4n-4i-2},\ M_n = v_n m_1' = M \prod_{i=0}^{n-1} A^{-1}_{4,4n-4i-2}.$$

□

7.2. **Proof of Theorem** 3.4. By Lemma 7.1, the dominant monomials in the q-characters of the right hand side and of the left hand side of every equation in Theorem 3.4 are the same. Therefore the theorem is true.

8. PROOF OF THEOREM 3.6

In this section, we prove Theorem 3.6.

By Lemma 7.1, we have known that the modules in the second summand on the right hand side of every equation in Theorem 3.4 are special, and hence they are simple. Therefore in order to prove Theorem 3.6, we only need to show that the modules in the first summand on the right hand side of every equation in Theorem 3.4 are simple.

8.1. **Proof of Theorem 3.6.** We will prove the case of $\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,k})$. The other cases are similar. By definition, we have

Equations	Non-highest dominant monomial	n_r	Relation
(3.1)	$M_r, \quad r=1,2,\ldots,k-1$	$M_rA^{-1}_{1,aq^{-2l-2r+1}}A^{-1}_{2,aq^{-2l-2r+2}}$, $r=1,2,\ldots,k-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\mathcal{T}^{(-2)}_{0,0,l-1,k})\chi_q(\mathcal{T}^{(-2)}_{0,0,l,k})$, $r=1,2,\ldots,k-1$
3.2)	$M_r, \quad r=1,2,\ldots,n-1$	$M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=1,2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,0,0})$, $r=1,2,\ldots,n-1$
(3.3, 3.4)	$M_r, \quad r=1,2,\ldots,n-1$	$M_1A^{-1}_{4,aq^{s+4n-2}}A^{-1}_{3,aq^{s+4n}}A^{-1}_{2,aq^{s+4n+2}}$ $M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,l-2,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,l,0})$, $r=1,2,\ldots,n-1$
(3.5, 3.6)	$M_r, \quad r=1,2,\ldots,m-1$	$M_rA^{-1}_{3,aq^{s+4m-4r+4}}A^{-1}_{2,aq^{s+4m-4r+6}}$, $r=1,2,\ldots,m-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,l-2,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,l,0})$, $r=1,2,\ldots,m-1$
(3.7)	$M_r, \quad r=1,2,\ldots,n-1$	$M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=1,2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,l,0})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,0})$, $r=1,2,\ldots,n-1$
(3.8)	$M_r, \quad r=1,2,\ldots,n-1$	$M_1A^{-1}_{4,aq^{s+4n-2}}A^{-1}_{3,aq^{s+4n}}A^{-1}_{2,aq^{s+4n+2}}A^{-1}_{1,aq^{s+4n+3}}$, $M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-2})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,0,k})$, $r=1,2,\ldots,n-1$
(3.9)	$M_r, \quad r=1,2,\ldots,m-1$	$M_1A^{-1}_{3,aq^{s+4m}}A^{-1}_{2,aq^{s+4m+2}}A^{-1}_{1,aq^{s+4m+3}}$, $M_rA^{-1}_{3,aq^{s+4m-4r+4}}A^{-1}_{2,aq^{s+4m-4r+6}}$, $r=2,\ldots,m-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,k-2})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,0,k})$, $r=1,2,\ldots,m-1$
(3.10, 3.11)	$M_r, \quad r=1,2,\ldots,n-1$	$M_1A^{-1}_{4,aq^{s+4n-2}}A^{-1}_{3,aq^{s+4n}}A^{-1}_{2,aq^{s+4n+2}}$, $M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-1})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,l,k})$, $r=1,2,\ldots,n-1$
(3.12, 3.13)	$M_r, \quad r=1,2,\ldots,m-1$	$M_rA^{-1}_{3,aq^{s+4m-4r+4}}A^{-1}_{2,aq^{s+4m-4r+6}}$, $r=1,2,\ldots,m-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,k-1})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{0,m,1,k})$, $r=1,2,\ldots,m-1$
(3.14)	$M_r, \quad r=1,2,\ldots,n-1$	$M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=1,2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,0,k})$, $r=1,2,\ldots,n-1$
(3.15)	$M_r, \quad r=1,2,\ldots,n-1$	$M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=1,2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k})$, $r=1,2,\ldots,n-1$
(3.16, 3.17, 3.18)	$M_r, \quad r=1,2,\ldots,l-1$	$M_rA^{-1}_{2,aq^{s+2l+4-2r}}A^{-1}_{1,aq^{s+2l+5-2r}}$, $r=1,2,\ldots,l-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{P}}^{(s+2)}_{0,0,l,k-1})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,0,l,k})$, $r=1,2,\ldots,l-1$
(3.19, 3.20)	$M_r, \quad r=1,2,\ldots,m-1$	$M_1A^{-1}_{3,aq^{s+4m}}A^{-1}_{2,aq^{s+4m+2}}A^{-1}_{1,aq^{s+4m+3}}$, $M_rA^{-1}_{3,aq^{s+4m-4r+4}}A^{-1}_{2,aq^{s+4m-4r+6}}$, $r=2,\ldots,m-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{P}}^{(s+4)}_{0,m,0,k-2})\chi_q(\widetilde{\mathcal{P}}^{(s)}_{0,m,0,k})$, $r=1,2,\ldots,m-1$
(3.21)	$M_r, \quad r=1,2,\ldots,n-1$	$M_1A^{-1}_{4,aq^{s+4n-2}}A^{-1}_{3,aq^{s+4n}}A^{-1}_{2,aq^{s+4n+2}}A^{-1}_{1,aq^{s+4n+3}}$, $M_rA^{-1}_{4,aq^{s+4n-4r+2}}A^{-1}_{3,aq^{s+4n-4r+4}}$, $r=2,\ldots,n-1$	$n_r\in\chi_q(M_r)$, $n_r\notin\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-2})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n,0,0,k})$, $r=1,2,\ldots,n-1$

TABLE 4. Irreducible in Theorem 3.4.

$$\begin{aligned}\widetilde{T}^{(s+4)}_{n,m-1,l,k} =&(4_{s+4}4_{4+8}\cdots 4_{s+4n})(3_{s+4n+6}3_{s+4n+10}\cdots 3_{s+4n+4m-2})(2_{s+4n+4m+3}2_{s+4n+4m+5}\cdots\\ &2_{s+4n+4m+2l+1})(1_{s+4n+4m+2l+4}1_{s+4n+4m+2l+6}\cdots 1_{s+4n+4m+2l+2k+2}),\\ \widetilde{T}^{(s)}_{n,m,l,k} =&(4_{s}4_{s+4}\cdots 4_{s+4n-4})(3_{s+4n+2}3_{s+4n+6}\cdots 3_{s+4n+4m-2})(2_{s+4n+4m+3}2_{s+4n+4m+5}\cdots\\ &2_{s+4n+4m+2l+1})(1_{s+4n+4m+2l+4}1_{s+4n+4m+2l+6}\cdots 1_{s+4n+4m+2l+2k+2}).\end{aligned}$$

By Lemma 7.1, the dominant monomials of $\chi_q(\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,k})\chi_q(\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,k})$ are

$$M_r = M\prod_{i=0}^{r-1} A^{-1}_{4,aq^{s+4n-2-4i}}, 0 \le r \le n-1, \text{ where } M = \widetilde{T}^{(s+4)}_{n-1,m,l,k}\widetilde{T}^{(s)}_{n+1,m-1,l,k}.$$

We need to show that $\chi_q(M_r) \not\subseteq \chi_q(M)$ for $1 \le r \le n-1$. We will prove the case of $r = 1$, the other cases are similar.

$$M_1 = MA^{-1}_{4,aq^{s+4n-2}} = M4^{-1}_{s+4n-4}4^{-1}_{s+4n}3_{s+4n-2}.$$

By $U_{q^2}(\widehat{\mathfrak{sl}}_2)$ argument, it is clear that $n_1 = M_1 A^{-1}_{4,aq^{s+4n-2}} A^{-1}_{3,aq^{s+4n}}$ is in $\chi_q(M_1)$.

If n_1 is in $\chi_q(\widetilde{T}^{(s+4)}_{n,m-1,l,k})\chi_q(\widetilde{T}^{(s)}_{n,m,l,k})$, then $\widetilde{T}^{(s+4)}_{n,m-1,l,k}A^{-1}_{4,aq^{s+4n-2}}A^{-1}_{3,aq^{s+4n}}$ is in $\chi_q(\widetilde{T}^{(s+4)}_{n,m-1,l,k})$ which is impossible by the Frenkel-Mukhin algorithm for $\widetilde{T}^{(s+4)}_{n,m-1,l,k}$. Hence $\chi_q(M_1) \not\subseteq \chi_q(M)$.

9. Conjectural equations satisfied by the q-characters of other minimal affinizations in type F_4

In this section, we give some conjectural equations satisfied by the q-characters of the minimal affinizations in type F_4 which are not in Theorem 3.4 and Theorem 5.4. In order to study equations satisfied by q-characters, we introduce the concept of dominant monomial graphs for a tensor product of simple $U_q\widehat{\mathfrak{g}}$-modules.

9.1. **Conjecture about the minimal affinizations which are not in Theorem 3.4 and Theorem 5.4.** Let $l, m, n \in \mathbb{Z}_{\geq 1}$, $k \in \mathbb{Z}_{\geq 3}$, $s \in \mathbb{Z}$. We define

$$\begin{aligned}
\widetilde{S}^{(s)}_{0,m,0,k} &= \widetilde{T}^{(s)}_{0,0,2m,k}\widetilde{T}^{(s+4)}_{m,0,0,k-2},\\
\widetilde{S}^{(s)}_{n,0,l,k} &= \begin{cases} \widetilde{T}^{(s)}_{0,n,1,k}\widetilde{T}^{(s+4n+4)}_{0,0,0,k-1}, & l = 1,\\ \widetilde{T}^{(s)}_{0,n,l,k}\widetilde{T}^{(s+4n+4)}_{0,0,l-2,k}, & l \geq 2,\end{cases}\\
\widetilde{S}^{(s)}_{0,m,l,k} &= \begin{cases} \widetilde{T}^{(s)}_{0,0,1+2m,k}\widetilde{T}^{(s+4)}_{m,0,0,k-1}, & l = 1,\\ \widetilde{T}^{(s)}_{0,0,l+2m,k}\widetilde{T}^{(s+4)}_{m,0,l-2,k}, & l \geq 2,\end{cases}\\
\widetilde{S}^{(s)}_{n,m,0,k} &= \widetilde{T}^{(s)}_{0,n+m,0,k}\widetilde{T}^{(s+4n+4)}_{0,m-1,0,k},\\
\widetilde{S}^{(s)}_{n,m,l,k} &= \widetilde{T}^{(s)}_{0,n+m,l,k}\widetilde{T}^{(s+4n+4)}_{0,m-1,l,k}.
\end{aligned}$$

We use $\widetilde{\mathcal{S}}^{(s)}_{n,m,l,k}$ to denote the simple $U_q\widehat{\mathfrak{g}}$-module with the highest weight monomial $\widetilde{S}^{(s)}_{n,m,l,k}$.

Conjecture 9.1. *For $s \in \mathbb{Z}$, $n, m, l \in \mathbb{Z}_{\geq 1}$, $k \in \mathbb{Z}_{\geq 3}$, we have the following equations in* $\operatorname{Rep}(U_q\widehat{\mathfrak{g}})$.

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,k-2}][\widetilde{\mathcal{T}}^{(s)}_{0,m,0,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,0,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,k-2}] + [\widetilde{\mathcal{S}}^{(s)}_{0,m,0,k}], \tag{9.1}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,0,k}][\widetilde{\mathcal{T}}^{(s)}_{n,m,0,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,0,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,0,k}] + [\widetilde{\mathcal{S}}^{(s)}_{n,m,0,k}], \tag{9.2}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,m-1,l,k}][\widetilde{\mathcal{T}}^{(s)}_{n,m,l,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,m,l,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,m-1,l,k}] + [\widetilde{\mathcal{S}}^{(s)}_{n,m,l,k}], \tag{9.3}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,0,k-1}][\widetilde{\mathcal{T}}^{(s)}_{n,0,1,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,1,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,0,k-1}] + [\widetilde{\mathcal{S}}^{(s)}_{n,0,1,k}], \quad l = 1, \tag{9.4}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{n,0,l-2,k}][\widetilde{\mathcal{T}}^{(s)}_{n,0,l,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{n-1,0,l,k}][\widetilde{\mathcal{T}}^{(s)}_{n+1,0,l-2,k}] + [\widetilde{\mathcal{S}}^{(s)}_{n,0,l,k}], \quad l \geq 2, \tag{9.5}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,0,k-1}][\widetilde{\mathcal{T}}^{(s)}_{0,m,1,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,1,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,0,k-1}] + [\widetilde{\mathcal{S}}^{(s)}_{0,m,1,k}], \quad l = 1, \tag{9.6}$$

$$[\widetilde{\mathcal{T}}^{(s+4)}_{0,m,l-2,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m,l,k}] = [\widetilde{\mathcal{T}}^{(s+4)}_{0,m-1,l,k}][\widetilde{\mathcal{T}}^{(s)}_{0,m+1,l-2,k}] + [\widetilde{\mathcal{S}}^{(s)}_{0,m,l,k}], \quad l \geq 2. \tag{9.7}$$

Example 9.2. *The following are some equations of Equation (9.1) in Conjecture 9.1.*

$$[1_0 3_{-6}][1_{-4}1_{-2}1_0 3_{-10}] = [1_{-4}1_{-2}1_0][1_0 3_{-10}3_{-6}] + [1_{-4}1_{-2}1_0^2 2_{-9}2_{-7}4_{-8}], \tag{9.8}$$

$$[1_0 3_{-10}3_{-6}][1_{-4}1_{-2}1_0 3_{-14}3_{-10}] = [1_{-4}1_{-2}1_0 3_{-10}][1_0 3_{-14}3_{-10}3_{-6}] + [1_{-4}1_{-2}1_0^2 2_{-13}2_{-11}2_{-9}2_{-7}4_{-12}4_{-8}], \tag{9.9}$$

$$[1_{-4}1_{-2}1_0 3_{-10}][1_{-8}1_{-6}1_{-4}1_{-2}1_0 3_{-14}] = [1_{-8}1_{-6}1_{-4}1_{-2}1_0][1_{-4}1_{-2}1_0 3_{-14}3_{-10}] + [1_{-8}1_{-6}1_{-4}^2 1_{-2}^2 1_0^2 2_{-13}2_{-11}4_{-12}]. \tag{9.10}$$

Example 9.3. *The following are some equations of Equation (9.4) and Equation (9.5) in Conjecture 9.1.*

$$[1_{-2}1_0 4_{-10}][1_{-4}1_{-2}1_0 2_{-7}4_{-14}] = [1_{-4}1_{-2}1_0 2_{-7}][1_{-2}1_0 4_{-14}4_{-10}] + [1_{-4}1_{-2}^2 1_0^2 2_{-7}3_{-12}], \tag{9.11}$$

$$[1_{-2}1_0 4_{-14}4_{-10}][1_{-4}1_{-2}1_0 2_{-7}4_{-18}4_{-14}] = [1_{-4}1_{-2}1_0 2_{-7}4_{-14}][1_{-2}1_0 4_{-18}4_{-14}4_{-10}] + [1_{-4}1_{-2}^2 1_0^2 2_{-7}3_{-16}3_{-12}], \tag{9.12}$$

$$[1_{-4}1_{-2}1_0 4_{-12}][1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-16}] = [1_{-4}1_{-2}1_0 2_{-9}2_{-7}][1_{-4}1_{-2}1_0 4_{-16}4_{-12}] + [1_{-4}^2 1_{-2}^2 1_0^2 2_{-9}2_{-7}3_{-14}], \tag{9.13}$$

$$\begin{aligned}[1_{-4}1_{-2}1_0 4_{-16}4_{-12}][1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-20}4_{-16}] &= [1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-16}][1_{-4}1_{-2}1_0 4_{-20}4_{-16}4_{-12}] \\ &\quad + [1_{-4}^2 1_{-2}^2 1_0^2 2_{-9}2_{-7}3_{-18}3_{-14}],\end{aligned} \tag{9.14}$$

$$\begin{aligned}[1_{-4}1_{-2}1_0 2_{-7}4_{-14}][1_{-4}1_{-2}1_0 2_{-11}2_{-9}2_{-7}4_{-18}] &= [1_{-4}1_{-2}1_0 2_{-11}2_{-9}2_{-7}][1_{-4}1_{-2}1_0 2_{-7}4_{-18}4_{-14}] \\ &\quad + [1_{-4}^2 1_{-2}^2 1_0^2 2_{-11}2_{-9}2_{-7}^2 3_{-16}],\end{aligned} \tag{9.15}$$

$$\begin{aligned}[1_{-4}1_{-2}1_0 2_{-7}4_{-18}4_{-14}][1_{-4}1_{-2}1_0 2_{-11}2_{-9}2_{-7}4_{-22}4_{-18}] &= [1_{-4}1_{-2}1_0 2_{-11}2_{-9}2_{-7}4_{-18}][1_{-4}1_{-2}1_0 2_{-7}4_{-22}4_{-18}4_{-14}] \\ &\quad + [1_{-4}^2 1_{-2}^2 1_0^2 2_{-11}2_{-9}2_{-7}^2 3_{-20}3_{-16}],\end{aligned} \tag{9.16}$$

$$\begin{aligned}[1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-16}][1_{-4}1_{-2}1_0 2_{-13}2_{-11}2_{-9}2_{-7}4_{-20}] &= [1_{-4}1_{-2}1_0 2_{-13}2_{-11}2_{-9}2_{-7}][1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-16}] \\ &\quad + [1_{-4}^2 1_{-2}^2 1_0^2 2_{-13}2_{-11}2_{-9}^2 2_{-7}^2 3_{-18}].\end{aligned} \tag{9.17}$$

9.2. **Dominant monomial graphs.** In order to study equations satisfied by q-characters, we introduce dominant monomial graphs for a tensor product of simple $U_q\widehat{\mathfrak{g}}$-modules.

Definition 9.4. *Let $\mathcal{T} = \mathcal{T}_1 \otimes \cdots \otimes \mathcal{T}_k$ be a tensor product of simple $U_q\widehat{\mathfrak{g}}$-modules. We define the dominant monomial graph $G(\mathcal{T})$ for $\mathcal{T}$ as follows. The vertices of $G(\mathcal{T})$ are dominant monomials in $\chi_q(\mathcal{T}) = \chi_q(\mathcal{T}_1) \cdots \chi_q(\mathcal{T}_k)$. For two vertices v_1, v_2 in $G(\mathcal{T})$, there is an arrow from v_1 to v_2 if and only if $v_2 < v_1$.*

Let G be a dominant monomial graph. Suppose that a, b are two vertices in G and $b < a$. Then $b = ma$ for some $m \in \mathcal{Q}^-$. We draw

$$a \xrightarrow{\quad m \quad} b$$

when we draw the graph G.

In the following, we draw the dominant monomial graphs for the modules in the equivalence classes on the left hand side of the equations in Examples 9.2 and 9.3. Figure 1 – Figure 10 correspond Equation (9.8) – Equation (9.17) respectively.

In all examples of dominant monomial graphs, we find that every graph can be divided into two parts. The vertices in the first (resp. second) part of the graph are dominant monomials in the first (resp. second) summand of the right hand side of the correpsonding equation. For example, in Figure 2, the monomials M, a_2, a_3, a_4 (resp. a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11}) are the dominant monomials of the first (resp. second) summand of the right hand side of Equation (9.9).

These graphs are also conjectural, since we are not able to show that the Frenkel-Mukhin algorithm works for the modules which are not special. If we can show that these graphs are indeed the dominant monomial graphs for the corresponding modules, then the corresponding conjectural equations are true.

For $k \in \mathbb{Z}$, let

$$\begin{aligned} N_k^{(1)} &= A_{4,k}^{-1} A_{3,k+2}^{-1} A_{2,k+4}^{-1} A_{1,k+5}^{-1}, \\ N_k^{(2)} &= A_{2,k}^{-1} A_{3,k+2}^{-1} A_{2,k+4}^{-1} A_{1,k+5}^{-1}, \\ N_k^{(3)} &= A_{4,k-2}^{-1} A_{3,k}^{-1}, \\ N_k^{(4)} &= A_{2,k-2}^{-1} A_{3,k}^{-1} A_{4,k-2}^{-1}, \\ M_k^{(1)} &= A_{2,k-2}^{-1} A_{3,k}^{-1} A_{4,k+2}^{-1}, \end{aligned}$$

where $A_{i,k} = A_{i,aq^k}$ which is defined in 2.2.

We have the following relations

$$\begin{aligned} A_{2,k} N_k^{(2)} = A_{4,k} N_k^{(1)}, \quad A_{2,k+2} A_{1,k+3} N_{k-2}^{(1)} = N_k^{(3)}, \\ N_k^{(3)} = A_{2,k-2} N_k^{(4)}, \quad A_{4,k-2} N_k^{(4)} = A_{4,k+2} M_k^{(1)}. \end{aligned}$$

$$\begin{tikzcd} M \arrow[d, "A_{3,-8}^{-1}"] \\ a_2 \arrow[d, "N_{-6}^{(2)} A_{4,-6}^{-1}"] \\ a_3 \arrow[d, "N_{-8}^{(2)}"] \\ a_4 \end{tikzcd}$$

FIGURE 1. The dominant monomial graph for $L(m_1) \bigotimes L(m_2)$ ($M = m_1 m_2$ where $m_1 = 1_0 3_{-6}, m_2 = 1_{-4} 1_{-2} 1_0 3_{-10}$)

.

$$\begin{tikzcd} M \arrow[r, "A_{3,-8}^{-1}"] & a_2 \arrow[r, "N_{-6}^{(2)} A_{4,-6}^{-1}"] \arrow[d, "A_{3,-12}^{-1}"] & a_3 \arrow[r, "N_{-8}^{(2)}"] \arrow[d, "A_{3,-12}^{-1}"] & a_4 \arrow[d, "A_{3,-12}^{-1}"] & & \\ & a_5 \arrow[r, "N_{-6}^{(2)} A_{4,-6}^{-1}"] & a_6 \arrow[r, "N_{-8}^{(2)}"] \arrow[d, "N_{-8}^{(4)} A_{2,-8}^{-1}"] & a_7 \arrow[d, "N_{-8}^{(4)}"] & & \\ & & a_8 \arrow[r, "N_{-8}^{(2)} A_{2,-8}"] & a_9 \arrow[r, "A_{2,-6}^{-1} A_{1,-5}^{-1}"] & a_{10} \arrow[r, "N_{-12}^{(2)}"] & a_{11} \end{tikzcd}$$

FIGURE 2. The dominant monomial graph for $L(m_1) \bigotimes L(m_2)$ ($M = m_1 m_2$ where $m_1 = 1_0 3_{-10} 3_{-6}, m_2 = 1_{-4} 1_{-2} 1_0 3_{-14} 3_{-10}$).

$$\begin{tikzcd}
M \arrow[r, "N^{(2)}_{-6}A^{-1}_{3,-8}A^{-1}_{4,-6}"] \arrow[d, "A^{-1}_{3,-12}"] & a_2 \arrow[r, "N^{(2)}_{-8}"] \arrow[d, "A^{-1}_{3,-12}A^{-1}_{2,-10}"] & a_3 \arrow[r, "N^{(2)}_{-10}A^{-1}_{3,-12}A^{-1}_{4,-10}"] \arrow[d, "A^{-1}_{3,-12}A^{-1}_{2,-10}M^{(1)}_{-10}"] & a_4 \arrow[r, "N^{(2)}_{-12}"] \arrow[ddl, "M^{(1)}_{-10}"] & a_5 \arrow[dd, "A^{-1}_{4,-8}"] \\
a_6 \arrow[r, "N^{(2)}_{-6}M^{(1)}_{-8}"] \arrow[d, "N^{(2)}_{-10}A^{-1}_{4,-10}"] & a_7 \arrow[r, "N^{(2)}_{-8}M^{(1)}_{-10}"] \arrow[d, "N^{(1)}_{-10}"] & a_8 \arrow[d, "N^{(1)}_{-10}"] & & \\
a_{11} \arrow[r, "N^{(2)}_{-6}A^{-1}_{3,-8}A^{-1}_{4,-6}"] \arrow[d, "N^{(2)}_{-12}"] & a_{12} \arrow[r, "N^{(2)}_{-8}M^{(1)}_{-10}"] \arrow[d, "N^{(2)}_{-12}"] & a_9 \arrow[rr, "A^{-1}_{2,-8}A^{-1}_{1,-7}"] & & a_{10} \\
a_{13} \arrow[r, "N^{(2)}_{-6}A^{-1}_{3,-8}A^{-1}_{4,-6}"] & a_{14} & & &
\end{tikzcd}$$

FIGURE 3. The dominant monomial graph for $L(m_1) \bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-4}1_{-2}1_0 3_{-10}, m_2 = 1_{-8}1_{-6}1_{-4}1_{-2}1_0 3_{-14}$).

$$\begin{tikzcd}
M \arrow[r, "N^{(2)}_{-6}"] \arrow[d, "A^{-1}_{4,-12}"] & a_2 \arrow[d, "A^{-1}_{4,-12}"] \\
a_3 \arrow[r, "N^{(2)}_{-6}"] & a_4
\end{tikzcd}$$

FIGURE 4. The dominant monomial graph for $L(m_1) \bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-2}1_0 4_{-10}, m_2 = 1_{-4}1_{-2}1_0 2_{-7}4_{-14}$).

$$\begin{tikzcd}
M \arrow[r, "N^{(2)}_{-6}"] \arrow[d, "A^{-1}_{4,-12}"] & a_2 \arrow[d, "A^{-1}_{4,-12}"] \\
a_3 \arrow[r, "N^{(2)}_{-6}"] \arrow[d, "A^{-1}_{4,-16}"] & a_4 \arrow[d, "A^{-1}_{4,-16}"] \\
a_5 \arrow[r, "N^{(2)}_{-6}"] & a_6
\end{tikzcd}$$

FIGURE 5. The dominant monomial graph for $L(m_1) \bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-2}1_0 4_{-14}4_{-10}, m_2 = 1_{-4}1_{-2}1_0 2_{-7}4_{-18}4_{-14}$).

$$\begin{tikzcd}
M \arrow[r, "N^{(2)}_{-6}"] \arrow[d, "A^{-1}_{4,-14}"] & a_2 \arrow[r, "N^{(2)}_{-8}"] \arrow[d, "A^{-1}_{4,-14}"] & a_3 \arrow[d, "A^{-1}_{4,-14}"] \\
a_4 \arrow[r, "N^{(2)}_{-6}"] & a_5 \arrow[r, "N^{(2)}_{-8}"] & a_6
\end{tikzcd}$$

FIGURE 6. The dominant monomial graph for $L(m_1) \bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-4}1_{-2}1_0 4_{-12}, m_1 = 1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-16}$).

$$\begin{tikzcd}
M \arrow[r, "N^{(2)}_{-6}"] \arrow[d, "A^{-1}_{4,-14}"'] & a_2 \arrow[r, "N^{(2)}_{-8}"] \arrow[d, "A^{-1}_{4,-14}"'] & a_3 \arrow[d, "A^{-1}_{4,-14}"'] \\
a_4 \arrow[r, "N^{(2)}_{-6}"] \arrow[d, "A^{-1}_{4,-18}"'] & a_5 \arrow[r, "N^{(2)}_{-8}"] \arrow[d, "A^{-1}_{4,-18}"'] & a_6 \arrow[d, "A^{-1}_{4,-18}"'] \\
a_7 \arrow[r, "N^{(2)}_{-6}"] & a_8 \arrow[r, "N^{(2)}_{-8}"] & a_9
\end{tikzcd}$$

FIGURE 7. The dominant monomial graph for $L(m_1)\bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-4}1_{-2}1_04_{-16}4_{-12}, m_2 = 1_{-4}1_{-2}1_02_{-9}2_{-7}4_{-20}4_{-16}$).



FIGURE 8. The dominant monomial graph for $L(m_1)\bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-4}1_{-2}1_02_{-7}4_{-14}, m_2 = 1_{-4}1_{-2}1_02_{-11}2_{-9}2_{-7}4_{-18}$).



FIGURE 9. The dominant monomial graph for $L(m_1)\bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-4}1_{-2}1_0 2_{-7}4_{-18}4_{-14}, m_2 = 1_{-4}1_{-2}1_0 2_{-11}2_{-9}2_{-7}4_{-22}4_{-18}$).



FIGURE 10. The dominant monomial graph for $L(m_1)\bigotimes L(m_2)$ ($M = m_1m_2$ where $m_1 = 1_{-4}1_{-2}1_0 2_{-9}2_{-7}4_{-16}, m_2 = 1_{-4}1_{-2}1_0 2_{-13}2_{-11}2_{-9}2_{-7}4_{-20}$).

ACKNOWLEDGEMENT

The authors are supported by the National Natural Science Foundation of China (no. 11371177, 11401275), the Fundamental Research Funds for the Central Universities of China.

References

[C95] V. Chari, *Minimal affinizations of representations of quantum groups: the rank* 2 *case*, Publ. Res. Inst. Math. Sci. **31** (1995), no. 5, 873–911.

[CG11] V. Chari, J. Greenstein, *Minimal affinizations as projective objects*, J. Geom. Phys. **61** (2011), no. 3, 594–609.

[CMY13] V. Chari, A. Moura, C. A. S. Young, *Prime representations from a homological perspective*, Math. Z. **274** (2013), 1-2, 613–645

[CP91] V. Chari, A. Pressley, *Quantum affine algebras*, Comm. Math. Phys. **142** (1991), no. 2, 261–283.

[CP94] V. Chari, A. Pressley, *A guide to quantum groups*, Cambridge University Press, Cambridge (1994), xvi+651 pp.

[CP95a] V. Chari, A. Pressley, *Quantum affine algebras and their representations*, Representations of groups (Banff, AB, 1994), 59–78, CMS Conf. Proc., **16**, Amer. Math. Soc., Providence, RI, 1995.

[CP96a] V. Chari, A. Pressley, *Minimal affinizations of representations of quantum groups: the irregular case*, Lett. Math. Phys. **36** (1996), no. 3, 247–266.

[Dri87] V. G. Drinfeld, *Quantum groups*, Proceedings of the International Congress of Mathematicians, Vol. **1**, **2** (Berkeley, Calif., 1986), Amer. Math. Soc., Providence, RI (1987), 798–820.

[Dri88] V. G. Drinfeld, *A new realization of Yangians and of quantum affine algebras*, (Russian) Dokl. Akad. Nauk SSSR **296** (1987), no. 1, 13–17; translation in Soviet Math. Dokl. **36** (1988), no. 2, 212–216.

[FM01] E. Frenkel, E. Mukhin, *Combinatorics of q-characters of finite-dimensional representations of quantum affine algebras*, Comm. Math. Phys. **216** (2001), no. 1, 23–57.

[FR98] E. Frenkel, N. Yu. Reshetikin, *The q-characters of representations of quantum affine algebras and deformations of W-algebras*, Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998), 163–205, Contemp. Math., **248**, Amer. Math. Soc., Providence, RI, 1999.

[FZ02] S. Fomin, A. Zelevinsky, *Cluster algebras I: Foundations*, J. Amer. math. Soc. **15** (2002), 497–529.

[GG14] J. Grabowski, S. Gratz, *Cluster algebras of infinite rank, with an appendix by Michael Groechenig*, J. Lond. Math. Soc. **89** (2014), no. 2, 337–363.

[Her06] D. Hernandez, *The Kirillov-Reshetikhin conjecture and solutions of T-systems*, J. Reine Angew. Math. **596** (2006), 63–87.

[Her07] D. Hernandez, *On minimal affinizations of representations of quantum groups*, Comm. Math. Phys. **276** (2007), no. 1, 221–259.

[HL10] D. Hernandez, B. Leclerc, *Cluster algebras and quantum affine algebras*, Duke Math. J. **154** (2010), no. 2, 265–341.

[HL13] D. Hernandez, B. Leclerc, *A cluster algebra approach to q-characters of Kirillov-Reshetikhin modules*, arXiv:1303.0744, 1–45.

[Jim85] M. Jimbo, *A q-difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), no. 1, 63–69.

[Li15] J. R. Li, *On the extended T-system of type C_3*, Journal of Algebraic Combinatorics, **41** (2015), 577-617.

[LM13] J. R. Li, E. Mukhin, *Extended T -system of type G_2*, SIGMA Symmetry Integrability Geom. Methods Appl. **9** (2013), 054, 28pp.

[M10] A. Moura, *Restricted limits of minimal affinizations*, Pacific J. Math. **244** (2010), no. 2, 359–397.

[MP07] M. G. Moakes, A. N. Pressley, *q-characters and minimal affinizations*, Int. Electron. J. Algebra **1** (2007), 55–97.

[MP11] A. Moura, F. Pereira, *Graded limits of minimal affinizations and beyond: the multiplicity free case for type E_6*, Algebra Discrete Math. **12** (2011), no. 1, 69–115.

[MY12a] E. Mukhin, C. A. S. Young, *Extended T-systems*, Selecta Math.(N.S.), **18** (2012), no. 3, 591–631.

[MY12b] E. Mukhin, C. A. S. Young, *Path description of type B q-characters*, Adv. Math. **231** (2012), no. 2, 1119–1150.

[MY14] E. Mukhin, C. A. S. Young, *Affinization of category $\mathcal{O}$ for quantum groups*, Trans. Amer. Math. Soc. **366** (2014), 4815–4847.

[Nao13] K. Naoi, *Demazure modules and graded limits of minimal affinizations*, Represent. Theory, **17** (2013), 524–556.

[Nao14] K. Naoi, *Graded limits of minimal affinizations in type D*, SIGMA Symmetry Integrability Geom. Methods Appl. **10** (2014), Paper 047, 20 pp.

[QL14] L. Qiao, J. R. Li, *Cluster algebras and minimal affinizations of representations of the quantum group of type* G_2, arXiv:1412.3884, 2014, 1–17.

[SS14] S. V. Sam, *Jacobi-Trudi determinants and characters of minimal affinizations*, Pacific J. Math. **272** (2014), no. 1, 237–244.

[ZDLL15] Q. Q. Zhang, B. Duan, J. R. Li, Y. F. Luo, *M-systems and cluster algebras*, arXiv:1501.00146, 2015, 1–44.

BING DUAN: SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU 730000, P. R. CHINA.
E-mail address: `duan890818@163.com`

JIAN-RONG LI: SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU 730000, P. R. CHINA.
E-mail address: `lijr@lzu.edu.cn`

YAN-FENG LUO: SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU 730000, P. R. CHINA.
E-mail address: `luoyf@lzu.edu.cn`